

HS-695T Cells | 300211**General information**

Description HS-695T was isolated in 1973 from the lymph node metastasis of an amelanotic melanoma by A. Creasey et al. The cells had a doubling time of approximately 48 hours at passages 19 through 40 and grew moderately in semi-solid medium.

Organism Human

Tissue Skin

Disease Amelanotic melanoma

Metastatic site Lymph node

Synonyms Hs 695.T, Hs-695-T, Hs 695T, HS 695T, Hs695T, HS695T, Hs695

Characteristics

Age 26 years

Gender Male

Ethnicity Caucasian

Morphology Epithelial-like

Growth properties Adherent

Identifiers / Biosafety / Citation

Citation HS-695T (Cytion catalog number 300211)

Biosafety level 1

Depositor R. B. Owens

Expression / Mutation

Protein expression p53 positive

HS-695T Cells | 300211

Isoenzymes	G6PD, B, PGM1, 1, PGM3, 1, ES-D, 1, Me-2, 0, AK-1, 1, GLO-1, 1, Phenotype Frequency Product: 0.0427
Tumorigenic	Yes, in immunosuppressed mice
Mutational profile	BRAF V600Emut
Karyotype	(P19-40) mode = 52, Y chromosome present

Handling

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Split ratio	A ratio of 1:2 to 1:4 is recommended
Seeding density	2 x 10 ⁴ cells/cm ²
Fluid renewal	2 to 3 times per week
Freezing recovery	After thawing, plate the cells at 5 x 10 ⁴ cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

HS-695T Cells | 300211

Handling of cryopreserved cultures

HS-695T cells are shipped in a deep-frozen state on dry ice. Upon receipt, confirm that the vial remains frozen. For storage, place the cryovial immediately at temperatures below -150 degrees. If you plan to culture the cells immediately, swiftly thaw the vial by shaking it in a 37 degrees water bath with clean water and an antimicrobial agent for 40-60 seconds. Remove the vial once a small ice clump persists, ensuring it remains cold. Proceed with all subsequent steps under aseptic conditions. In a sterile flow hood, disinfect the cryovial with 70% ethanol. Then, gently open the vial and transfer the cell suspension into a 15 ml centrifuge tube pre-filled with 8 ml of room temperature culture medium. Gently mix the cells. For cell separation, centrifuge at 300 x g for 3 minutes and dispose of the supernatant. Skipping centrifugation is optional, although any residual freezing medium should be removed after 24 hours. Resuspend the pellet gently in 10 ml of fresh culture medium and divide between two T25 culture flasks. Follow the subculture protocol for subsequent steps.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is rigorously excluded using both PCR-based assays and luminescence-based mycoplasma detection methods. To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

Amelogenin: x,y
CSF1PO: 11
D13S317: 12
D16S539: 9,13
D5S818: 9
D7S820: 9,10
TH01: 6
TPOX: 8
vWA: 18
D3S1358: 15
D21S11: 29
D18S51: 18
Penta E: 5,11
Penta D: 9,12
D8S1179: 13,15
FGA: 21,24