

INS-1 Cells | 300471

General information

Description

The INS-1 cell line is derived from an x-ray-created transplantable insulinoma in rats. Because INS-1 cells contain a high concentration of insulin and respond to changes in glucose levels, they are frequently used to study the function of beta cells. Growth and hormone expression are dependent on the reducing agent 2-mercaptoethanol.

INS-1 cells are notable for their heterogeneity, consisting of mature insulin-positive cells and immature bi-hormonal cells expressing insulin and glucagon proteins.

Bi-hormonal INS-1 cells have lower Nkx6.1 expression and lack alpha cell markers, indicating they are not fully matured. Furthermore, chronic glucose stimulation reduces insulin gene and protein expression in INS-1 cells. As such, insulin and proglucagon-derived peptides like GLP-1, GLP-2, and glucagon levels are reduced.

Organism Rat

Tissue Pancreas, islets of Langerhans

Disease Rat insulinoma

Synonyms INS1

Characteristics

Breed/Subspecies NEDH

Age 666 days

Gender Male

Cell type Beta cell

Growth properties Adherent/suspension

Regulatory Data

Citation INS-1 (Cytion catalog number 300471)

Biosafety level 1

INS-1 Cells | 300471**NCBI_TaxID** 10116**CellosaurusAccession** CVCL_0352**Biomolecular Data****Products** Insulin, glutathione**Handling****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)**Supplements** Supplement the medium with 10% heat-inactivated FBS, add 2.5 g/L glucose and 10 mM HEPES**Dissociation Reagent** Accutase**Subculturing** Gather the suspension cells in a 15 ml tube and gently wash the adherent cells with PBS lacking calcium and magnesium (use 3-5 ml for T25 flasks and 5-10 ml for T75 flasks). Apply Accutase (1-2 ml for T25 flasks, 2.5 ml for T75 flasks) ensuring full coverage of the cell layer. Allow the cells to incubate at room temperature for 10 minutes. Following incubation, combine and centrifuge both the suspension and adherent cells. After centrifugation, carefully resuspend the cell pellet and transfer the cell suspension into new flasks containing fresh medium.**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

INS-1 Cells | 300471

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Flask Coating

None

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

INS-1 Cells | 300471

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.