

RAJI Cells | 300359**General information****Description**

Raji cells are a line of lymphoblast-like cells established by R.J.V. Pulvertaft in 1963 from Burkitt's lymphoma. These cells are widely used in immunology research due to their high expression of human CD19, which acts as a co-receptor and decreases the threshold for antigen B-cell receptor (BCR) stimulation. Raji cells are non-adherent and grow in suspension as free-floating individuals or doublets.

The doubling time of these cells is 23.2 hours, and they are relatively small in diameter with a diameter range of 5-8 µm. Some characteristics of Raji cells include a lack of differentiation, as they form large aggregations of hundreds of individual cells. These cells are diploid and have a stable karyotype within the male diploid stemline of 46.

Additionally, Raji cells are partially resistant to poliovirus and vesicular stomatitis viruses. Human CD19 is highly expressed by Raji cells and has been identified as a clinical target for anti-hCD19-CD3 bis-specific antibodies in non-Hodgkin's B cell lymphoma. BCMA expression has also been identified in the Raji Burkitt lymphoma cell line and primary lymphoma, making it an important area of research for immunologists.

Organism

Human

Tissue

Maxilia

Disease

Burkitt lymphoma

Synonyms

Raji, P1-Raji, GM04671

Characteristics**Age**

11 years

Gender

Male

Ethnicity

African, Nigerian

Cell type

Lymphoblast

Growth properties

Suspension

Identifiers / Biosafety / Citation**Citation**

RAJI (Cytion catalog number 300359)

RAJI Cells | 300359

Biosafety level 1

Expression / Mutation

Products	The cells may produce interferon when stimulated by Newcastle disease virus.
-----------------	--

Handling

Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Subculturing	Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of 1×10^5 cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation.
---------------------	--

Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
----------------------	--

RAJI Cells | 300359

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

RAJI Cells | 300359

STR profile

CSF1PO: 10,12
D13S317: 13
D16S539: 8,11
D5S818: 10,13
D7S820: 10
TH01: 6,7
TPOX: 8,13
vWA: 16,19
D3S1358: 15,16
D21S11: 28,31
D18S51: 17
Penta E: 5,13
Penta D: 3,2,9
D8S1179: 14,15
FGA: 19,27

HLA alleles

A*: 03:01:01
B*: 15:10:01
C*: 03:04:02, 04:01:01
DRB1*: 03:01:01, 10:01:01
DQA1*: 01:05:01, 05:01:01
DQB1*: 02:01:01, 05:01:01
DPB1*: 01:01:01
E: 01:01:01