

General information

Description

AGS cells are a human gastric adenocarcinoma cell line derived from the stomach tissue of a 54-year-old Caucasian female. They are extensively used in biomedical research focused on gastric cancer, including studies on cancer cell biology, pathogenesis, and drug testing.

The AGS cell line exhibits epithelial-like morphology and is characterized by its aggressive growth pattern and tumorigenic potential in vivo. These cells are commonly used as a model to study the molecular and cellular mechanisms underlying gastric carcinogenesis, including the influence of *Helicobacter pylori* infection, a well-known risk factor for gastric cancer. AGS cells provide a robust system to explore the interactions between gastric cancer cells and *H. pylori*, especially regarding how bacterial factors affect cancer cell proliferation, apoptosis, and inflammatory responses.

AGS cells are also valuable for examining the gastric epithelial barrier's response to various stimuli, including inflammatory cytokines, and for studying signaling pathways implicated in gastric cancer, such as those involving NF- κ B, Wnt, and MAPK. Their utility extends to the assessment of new therapeutic agents, where they are used to evaluate the efficacy and mechanisms of action of anticancer drugs, targeted therapies, and natural compounds with potential anti-cancer properties.

Furthermore, AGS cells are often employed in studies aimed at understanding the genetic and epigenetic alterations in gastric cancer, offering insights into potential diagnostic markers and therapeutic targets for this challenging and frequently fatal disease.

Organism

Human

Tissue

Gastric

Disease

Adenocarcinoma

Characteristics

Age

54 years

Gender

Female

Ethnicity

Caucasian

Morphology

Epithelial-like

Growth properties

Monolayer, adherent

Identifiers / Biosafety / Citation

AGS Cells | 300408

Citation AGS (Cyton catalog number 300408)

Biosafety level 2

Expression / Mutation

Protein expression p53 positive

Tumorigenic Yes, in athymic BALB/c mice

Viruses This cell line may release Parainfluenzavirus Type 5 (formerly known as Simian Virus 5). The virus interferes with Interferon-signalling within the cell line by degradation of STAT1.

Karyotype Modal number = 47, range = 39 to 92

Handling

Culture Medium DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO₃, w: 1.0 mM Sodium pyruvate (Cyton article number 820300a)

Medium supplements Supplement the medium with 10% FBS

Passaging solution Accutase

Doubling time 24 to 48 hours

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Split ratio A ratio of 1:2 to 1:6 is recommended

Seeding density 1×10^4 cells/cm² will result in a confluent monolayer within 3 to 5 days.

Fluid renewal 2 to 3 times per week

AGS Cells | 300408

Freeze medium

CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

AGS Cells | 300408

STR profile

Amelogenin: x,x
CSF1PO: 11,12
D13S317: 12
D16S539: 11,13
D5S818: 9,12
D7S820: 10,11
TH01: 6,7
TPOX: 11,12
vWA: 16,17
D3S1358: 16
D21S11: 29
D18S51: 13
Penta E: 13,16
Penta D: 9,10
D8S1179: 13
FGA: 23,24

HLA alleles

A*: 02:01:01
B*: 52:01:02
C*: 07:02:01
DRB1*: 08:02:01
DQA1*: 04:01:01
DQB1*: 04:02:01
DPB1*: 02:01:02
E: 01:03:02