

General information

Description	THP1 cells, a spontaneously immortalized monocyte-like cell line derived from the peripheral blood of a 1-year- old monocytic leukemia patient, serve as a critical model in immunological and cancer research. The THP-1 monocyte cell line, known for its ability to differentiate into mature macrophage and dendritic cells, is essential for studying the functions and properties of these immune cells in vitro, including adipose tissue macrophages and M2 mononuclear phagocytes.
	THP-1 differentiated macrophages are instrumental in exploring monocyte and macrophage functions, mechanisms, signalling pathways, including cytokine activation and immune modulation, and studying nutrient and drug transport. Further, THP-1 macrophages can be polarized into M1 or M2 macrophages, crucial for studies on immunity and inflammation, innate immunity, and inflammatory responses.
	In the context of metabolic and inflammatory diseases, THP-1 cells help explore cytokine profiles, including inflammatory cytokines, and their impact on conditions like human adipocyte apoptosis, illustrating the interplay between inflammation and metabolic health.
	Notably, the THP-1 cell line allows for comparative studies with other monocytic leukemia cells and cell lines like U937, facilitating a deeper understanding of monocyte and macrophage biology across different models.
	In summary, the THP-1 human monocytic leukemia cell line stands as a valuable tool for a myriad of research avenues, from investigating the intricate mechanisms of the immune system and its role in cancer to understanding the cellular and molecular underpinnings of immune modulation, cytokine activation, and cell proliferation. Its ability to mimic human macrophages and dendritic cells, combined with the ease of manipulation and fast growth rate, cements its status as a widely used cell line in biological and medical research, offering insights into the cellular basis of immunity and inflammation, the response of cancer cells, and the potential for therapeutic intervention.

Organism	Human
Tissue	The tissue of origin is peripheral blood
Disease	Leukemia
Applications	THP1 cells are a multifacted model with applications in Immune response modeling, Monocyte/macrophage differentiation, Phagocytosis mechanisms, Inflammatory signaling pathways, Drug transport assays
Synonyms	THP1, THP 1, THPI, O-THP-1, Tohoku Hospital Pediatrics-1

Characteristics

Age	1 year
Gender	Male



Morphology	Round cells
Cell type	Monocyte
Growth properties	The monocytic leukemia THP1 cell line grows in suspension and forms clumps due to cells dividing and attaching to the clusters they split from.

Identifiers / Biosafety / Citation

Citation	THP-1 (Cytion catalog number 300356)
Biosafety level	1

Expression / Mutation

Receptors expressed	HLA haplotypes: HLA-A2, -A9, -B5, -DRw1, -DRw2Fc, C3b
lsoenzymes	The human THP-1 cell line expresses low levels of CD4, CCR5, and CxCR4, making it relevant to HIV infection studies. However, they express low levels of CD14 and not CD80, CD86, CD11b, CD11c, Mertk, or CD1a, making them a poor model for primary monocytes regarding LPS responses.
Products	Lysozyme
Karyotype	THP-1 cells are near-diploid and contain two related subclones with genetic aberrations.
Handling	
Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)
Medium supplements	Supplement the medium with 10% FBS
Doubling time	The population doubling time of human THP-1 cells ranges from 19 to 50 hours, with an average of around 35 hours.
Subculturing	Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of 1 x 10^5 cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation.
Split ratio	Start cultures at 1 x 10^5 cells/ml and do not allow the cell concentration to exceed 1 x 10^6 cells/ml



Fluid renewal	2 to 3 times per week
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
Handling of cryopreserved cultures	1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
	2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
	3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
	4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
	5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
	6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
	7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
	8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.
Quality contro	l / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.



STR profile	CSF1PO: 11,13 D13S317: 13 D16S539: 11,12 D5S818: 11,12 D7S820: 10 TH01: 8,9.3 TPOX: 8,11 vWA: 16 D3S1358: 15,17 D21S11: 30,31.2 D18S51: 13,14 Penta E: 11,15 Penta D: 10,12 D8S1179: 10,14 FGA: 24,25
HLA alleles	A*: 02:01:01 B*: 15:11:01 C*: 03:03:01 DRB1*: 01:01:01, 15:01:01 DQA1*: 01:01:01, 01:02:01 DQB1*: 05:01:01, 06:02:01 DPB1*: 02:01:02G, 04:02:01G E: 01:03:02