

**SW620 Cells | 300466**

**General information**

**Description**

The SW-620 cell line, originating from the large intestine of a 51-year-old male with Dukes-C colorectal cancer, is a pivotal research model in colorectal cancer, especially for cancer biomarkers, chemotherapy, and the study of metastatic cancer cells.

SW-620 cells are pivotal for studying cell apoptosis and the resistance mechanisms to anoikis, a form of programmed cell death crucial for preventing metastasis. Research utilizing the SW-620 colon cancer cells has delved into proteomic analysis to understand proteome changes under different conditions, such as hypoxia. Hypoxic SW620 cells exhibit specific proteome adaptations that contribute to chemotherapy resistance.

SW620 colon cancer cells have been pivotal in evaluating natural compounds like curcumin and their impact on cancer cell viability. Studies have shown that curcumin inhibits cell viability in SW-620 cells. Moreover, the cell line aids in assessing the effects of chemotherapeutic agents and the potential for chemotherapy resistance, which is critical for advancing cancer treatment strategies.

Exhibiting high tumorigenic and metastatic capabilities, SW-620 cells form solid tumors in vivo. The SW620 xenograft model, alongside the study of specific pathways like the catenin pathway and the role of transcription factors such as cdx2 in colonic adenocarcinoma cells, enriches our understanding of colorectal cancer's molecular underpinnings.

In summary, SW-620 human colon adenocarcinoma cells are an invaluable resource in cancer research, offering a multifaceted approach to understanding colorectal cancer's complexities.

**Organism**

Human

**Tissue**

Colorectal

**Disease**

Adenocarcinoma

**Metastatic site**

Lymph node

**Synonyms**

SW620, SW 620, SW.620

**Characteristics**

**Age**

51 years

**Gender**

Male

**Ethnicity**

Caucasian

**Morphology**

Epithelial-like

## SW620 Cells | 300466

<b>Growth properties</b>	Adherent
--------------------------	----------

## Regulatory Data

<b>Citation</b>	SW-620 (Cytion catalog number 300466)
-----------------	---------------------------------------

<b>Biosafety level</b>	1
------------------------	---

<b>NCBI_TaxID</b>	9606
-------------------	------

<b>CellosaurusAccession</b>	CVCL_0547
-----------------------------	-----------

## Biomolecular Data

<b>Tumorigenic</b>	Yes, in athymic nude mice
--------------------	---------------------------

<b>Karyotype</b>	Average number of chromosomes 48 (range, 46-52). Eighteen marker chromosomes. For a detailed description of the karyotype we refer to Melcher et al.
------------------	--

## Handling

<b>Culture Medium</b>	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO <sub>3</sub> , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

<b>Supplements</b>	Supplement the medium with 10% FBS
--------------------	------------------------------------

<b>Dissociation Reagent</b>	Accutase
-----------------------------	----------

<b>Subculturing</b>	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

<b>Fluid renewal</b>	2 times per week
----------------------	------------------

<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	---

## SW620 Cells | 300466

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

**SW620 Cells | 300466**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.