

HK-2xZFN-Smc4-mEGFP Cells | 301576

General information

Description

The HK-2xZFN-Smc4-mEGFP cell line is derived from HeLa Kyoto cells and engineered using Zinc Finger Nuclease (ZFN) technology to target the Smc4 gene. Smc4 is crucial for chromosome condensation and segregation during cell division. This modification helps researchers study chromosomal dynamics and stability.

This cell line also features the monomeric Enhanced Green Fluorescent Protein (mEGFP) tag, allowing for real-time visualization of cellular processes. The mEGFP tag facilitates the observation of Smc4 expression and localization, aiding in high-throughput screening and live-cell imaging. The HK-2xZFN-Smc4-mEGFP cell line is a valuable tool for studying chromosomal behavior and gene function.

Organism Human

Tissue Endocervix

Disease Adenocarcinoma

Characteristics

Age 30 years

Gender Female

Ethnicity African American

Morphology Epithelial-like cells with mosaic stone shape

Growth properties Adherent

Regulatory Data

Citation HK-2xZFN-Smc4-mEGFP (Cytion catalog number 301576)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_B7FT

Depositor The Ellenberg Lab (EMBL)

HK-2xZFN-Smc4-mEGFP Cells | 301576

GMO Status GMO-S1: This HeLa Kyoto line contains a ZFN-engineered mEGFP tag on Smc4 for studying chromosome condensation dynamics. This classification applies only within Germany and may differ elsewhere.

Biomolecular Data

Products EGFP (Enhanced Green Fluorescent Protein)

Handling

Culture Medium DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO₃, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Fluid renewal 2 to 3 times per week

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HK-2xZFN-Smc4-mEGFP Cells | 301576

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HK-2xZFN-Smc4-mEGFP Cells | 301576

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.