



General information

DescriptionEstablished from the primary bladder carcinoma of a 1 year-old male ACI rat.OrganismRatTissueBladderDiseaseCarcinomaSynonymsHBCLS3

Characteristics

Age1 yearGenderMaleMorphologyEpithelial-likeGrowth propertiesAdherent

Identifiers / Biosafety / Citation

Citation HB-CLS-3 (Cytion catalog number 500460)

Expression / Mutation

Biosafety level

Tumorigenic Yes, in ACI-rats

Handling

 Culture Medium
 RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)

 Medium supplements
 Supplement the medium with 10% FBS



HB-CLS-3 Cells | 500460

| Passaging solution | Accutase |
|----------------------|---|
| Subculturing | Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium. |
| Split ratio | A ratio of 1:4 to 1:8 is recommended |
| Seeding density | 1 x 10^4 cells/cm^2 |
| Fluid renewal | 2 to 3 times per week |
| Freezing recovery | After thawing, plate the cells at 5×10^4 cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours. |
| Freeze medium | CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100) |



HB-CLS-3 Cells | 500460

Handling of cryopreserved cultures

- 1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
- 2. Upon receipt, either store the cryovial immediately at temperatures below -150?C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
- 3. For immediate culturing, swiftly thaw the vial by immersing it in a 37?C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
- 4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
- 5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
- 6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
- 7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
- 8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

HB-CLS-3 Cells | 500460



STR profile Rat_D1Wox31: 100

Rat_D2Wox37: 156
Rat_D19Wox11: 228
Rat_D10Wox8: 266,270
Rat_D4Wox7: 141,145
Rat_D2Wox27: 223
Rat_D5Rat33: 116,120,122
Rat_D10Wox11: 156,159
Rat_D1Wox23: 226,234

Rat_D12Wox1: 410 Rat_D6Wox2: 100,112,120 Rat_D8Wox7: 161,182 Rat_D6Cebr1: 239,241

SRY: x,x