

**GIMEN Cells | 300179****General information****Description**

GIMEN cells represent a distinct line of genetically engineered mouse embryonic neurons, meticulously developed to support advanced research in neurobiology and genetic disorders. These cells are derived from mouse embryonic stem cells and are modified to exhibit specific neuronal properties, making them highly suitable for studies on neural development, neurodegenerative diseases, and neuronal function.

The GIMEN cell line is characterized by its stable expression of both wild-type and mutant forms of human genes implicated in neurological conditions, allowing researchers to dissect the molecular pathways involved in disease states. These cells are cultured under strictly controlled conditions to maintain their undifferentiated state and neuronal potential. Upon differentiation, GIMEN cells are capable of expressing a variety of neuronal markers, including NeuN, MAP2, and beta-III tubulin, indicating their utility in modeling complex neural behaviors.

Furthermore, the GIMEN cells are amenable to a range of experimental manipulations, including gene editing, drug screening, and electrophysiological assessment, facilitating a comprehensive approach to neurological research. Their consistent performance and high genetic fidelity make them a valuable resource for both fundamental neuroscience research and the exploration of therapeutic interventions.

**Organism**

Human

**Tissue**

Brain

**Disease**

Neuroblastoma

**Metastatic site**

Bone marrow

**Synonyms**

Gi-ME-N, Gi-MEN, GI-ME-N, Gimen, Gimen1, Gaslini Institute-ME-Neuroblastoma

**Characteristics****Age**

3,5 years

**Gender**

Female

**Ethnicity**

Caucasian

**Morphology**

Epithelial-like

**Growth properties**

Adherent

## GIMEN Cells | 300179

### Identifiers / Biosafety / Citation

**Citation** GIMEN (Cytion catalog number 300179)

**Biosafety level** 1

### Expression / Mutation

### Handling

**Culture Medium** DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO<sub>3</sub>, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

**Medium supplements** Supplement the medium with 10% FBS

**Passaging solution** Accutase

**Doubling time** 25 hours

**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

**Seeding density** 2 to 3 x 10<sup>4</sup> cells/cm<sup>2</sup>

**Fluid renewal** 2 to 3 times per week

**Freeze medium** CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

### GIMEN Cells | 300179

#### Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Quality control / Genetic profile / HLA

#### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

## GIMEN Cells | 300179

### STR profile

**Amelogenin:** x,x  
**CSF1PO:** 11,12  
**D13S317:** 12  
**D16S539:** 9,12  
**D5S818:** 12  
**D7S820:** 10,11  
**TH01:** 6,7  
**TPOX:** 11  
**vWA:** 16,19  
**D3S1358:** 14  
**FGA:** 31  
**D1S1656:** 12,17  
**D6S1043:** 15,2  
**D2S1338:** 9,13  
**D12S391:** 10,14  
**D19S433:** 19,22

### HLA alleles

**A\*:** 02:01:01, 30:01:01  
**B\*:** 13:02:01, 18:01:01  
**C\*:** 06:02:01, 07:01:09  
**DRB1\*:** 04:03:01, 07:01:01  
**DQA1\*:** 02:01:01, 03:01:01  
**DQB1\*:** 02:02:01, 03:02:01  
**DPB1\*:** 02:01:02  
**E:** 01:01:01, 01:xx