

HMV-1 Cells | 305145

General information

Description

The HMV-1 cell line is a malignant melanoma cell line derived from a human male patient. The melanoma originated on the right sole of the patient and later metastasized to the lymph nodes. The cells exhibit a fibroblast-like morphology, characteristic of melanoma cells adapted for in vitro culture. Initially, the cell line exhibited pigmentation, a common trait in melanomas, but this pigmentation faded as the cell line became established and stabilized over time.

Genetically, the HMV-1 cell line has a modal chromosome number of 66, indicating significant chromosomal abnormalities typical of cancer cells. The cell line demonstrates an infinite lifespan in culture, providing a valuable model for long-term studies on melanoma biology, metastasis, and therapeutic interventions. As with other melanoma cell lines, HMV-1 can be utilized for research into the mechanisms of melanoma progression, drug resistance, and the development of novel treatments targeting metastatic cancer cells.

Organism Human

Tissue Skin

Disease Melanoma

Metastatic site Left inguinal lymph node

Synonyms HMV1

Characteristics

Age 62 years

Gender Male

Morphology Fibroblast

Growth properties Adherent

Regulatory Data

Citation HMV-1 (Cytion catalog number 305145)

Biosafety level 1

HMY-1 Cells | 305145

NCBI_TaxID 9606**CellosaurusAccession** CVCL_2950**Biomolecular Data****Tumorigenic** Yes**Handling****Culture Medium** DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO₃, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)**Supplements** Supplement the medium with 10% FBS**Dissociation Reagent** Accutase**Doubling time** 37 hours**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HMY-1 Cells | 305145

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HMY-1 Cells | 305145

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.