

## **General information**

Organism	Human
Tissue	Breast
Disease	Breast carcinoma
Metastatic site	Skin
Applications	3D cell culture, Immuno-oncology
Synonyms	Du4475, DU-4475, Du-4475, DU 4475, Du 4475, Duke University 4475

## Characteristics

Age	62 years
Gender	Female
Ethnicity	European
Morphology	Epithelial
Growth properties	Clusters in Suspension

# Identifiers / Biosafety / Citation

Citation	DU4475 (Cytion catalog number 300371)
<b>Biosafety level</b>	1

## **Expression / Mutation**

lsoenzymes	AK-1, 1, ES-D, 1, G6PD, B, GLO-I, 2, Me-2, 2, PGM1, 1-2, PGM3, 1
Tumorigenic	Yes, in nude mice
Viruses	EBV -, HBV -, HCV -, HIV-1 -, HIV-2 -, HTLV-1/2 -, MLV -, SMRV -

CLS Cell Lines Service GmbH | Dr.-Eckener-Str. 8 | 69214 Eppelheim | Germany Tel.: +49(0)6221 405780 | www.cytion.com | info@cytion.com



Karyotype Handling	human flat-moded near-tetraploid karyotype with 12% polyploidy - 88-93xxxx, +1, +1, -5, -6, +9, -10, -10, +15, +15, -16, -16, +22, +4mar, i(1q)x2, ?add(1)(p35-36)x2, ?i(5p)x2, add(6)(p11), add(6)(p1?), del(6)(q25), add(9)(q35), del(11)(q24)x2, add(15)(p11)x2, add(17)(p1?)x2, del(21)(q22.2)x2 - sideline with -20, -20, +del(7)(p11) - gain of 1q and loss of 6q typical in breast carcinoma - resembles published karyotype
nanating	
Culture Medium	RPMI 1640, w: 4.5 g/L Glucose, w: 2 mM L-Glutamine, w: 10 mM HEPES, w: 1 mM Sodium pyruvate, w: 1.5 g/L NaHCO3 (Cytion article number 820702a)
Medium supplements	Supplement the medium with 10% heat-inactivated FBS
Subculturing	Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 2 x 10^5 cells/ml and keep the cell concentration within the range of 1 x 10^5 to 1 x 10^6 cells/ml for optimal growth.
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)



Handling of cryopreserved cultures	1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
	2. Upon receipt, either store the cryovial immediately at temperatures below -150?C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
	3. For immediate culturing, swiftly thaw the vial by immersing it in a 37?C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
	4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
	5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
	6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
	<ol> <li>Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.</li> </ol>
	8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

# Quality control / Genetic profile / HLA

#### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.



Amelogenin: x,x CSF1PO: 9,12 D13S317: 11,14 D16S539: 11,12 D5S818: 11 D7S820: 9,10 **TH01**: 6,8 **TPOX**: 8 **vWA**: 17 D3S1358: 14,16 **D21S11**: 29,31.2 **D18S51**: 14,16 Penta E: 7,13 Penta D: 13,14 D8S1179: 10,13 FGA: 22,25 D6S1043: 11 **D2S1338**: 20,25 D12S391: 18.3,25 **D19S433**: 14