

HROHep03 Cells | 300197

General information

Description This is one cell line of a series of tumor cell lines which have been established by PD Dr. Michael Linnebacher

since 2006.

Organism Human

Tissue Liver

Disease Primary adenocarcinoma, T0NxMx stage, grade 3

Characteristics

Age 71 years

Gender Female

Ethnicity Caucasian

Morphology Fibroblast-like

Growth properties

Adherent

Identifiers / Biosafety / Citation

Citation HROHep03 (Cytion catalog number 300197)

Biosafety level 1

Depositor M. Linnebacher

Expression / Mutation

Viruses Free of human pathogenic viruses HBV, HCV, HIV.

Handling

CultureDMEM:Ham's F12, w: 3.1 g/L Glucose, w: 1.6 mM L-Glutamine, w: 15 mM HEPES, w: 1.0 mM Sodium pyruvate, w:

Medium
1.2 g/L NaHCO3 (Cytion article number 820400a)



HROHep03 Cells | 300197

Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	2 x 10^4 cells/cm^2
Fluid renewal	Every 3 to 5 days
Freezing recovery	2 days
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)



HROHep03 Cells | 300197

Handling of cryopreserved cultures

- 1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
- 2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
- 3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
- 4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
- 5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
- 6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
- 7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
- 8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.



HROHep03 Cells | 300197

STR profile Amelogenin: x,x

CSF1PO: 9,11 D13S317: 11,12 D16S539: 9,12 D5S818: 10,12 D7S820: 8,11 THO1: 7,9 TPOX: 8 vWA: 16,17 D3S1358: 15,16 D21S11: 30 D18S51: 12,17 D8S1179: 13,14 FGA: 19,22 D2S1338: 18,19 D19S433: 14,14.2