

HROC348Met Cells | 300871

General information

Description

HROC348Met is a human colorectal carcinoma cell line established from a metachronous liver metastasis of a colorectal adenocarcinoma resected from an adult patient within the HROC (Hansestadt Rostock Colorectal Cancer) model collection. The HROC platform was generated through a standardized biobanking and tumor-modeling pipeline integrating clinical annotation, molecular characterization, patient-derived xenografts (PDX), and corresponding in vitro cultures. HROC348Met represents one of the metastatic models derived from surgically resected colorectal cancer tissue and was established under low-passage conditions to preserve tumor-specific biological features.

Within the HROC collection, metastatic specimens - particularly liver metastases - demonstrated a high engraftment efficiency in immunodeficient mice, with an overall PDX take rate of approximately 68% across the cohort, and even higher success for metastatic compared to primary tumors. Multivariate analyses identified nodal involvement and activating mutations in KRAS and BRAF as independent predictors of successful model establishment. The collection encompasses all major molecular subtypes of colorectal carcinoma, including chromosomal instability (CIN), CpG island methylator phenotype (CIMP), microsatellite stable (MSS), and microsatellite instability-high (MSI-H) tumors, ensuring molecular representativeness of advanced-stage disease. HROC348Met was established within this rigorously characterized framework, with clinicopathological and molecular annotation according to standardized protocols.

As a metastasis-derived, low-passage colorectal carcinoma model, HROC348Met is suitable for investigations of metastatic tumor biology, genotype-phenotype correlations, and therapeutic response testing in both 2D culture and in vivo PDX settings. The integrated biobank approach underlying its generation ensures availability of matched clinical data and, where applicable, corresponding xenograft material, enabling translational studies in precision oncology and drug-response prediction.

Organism Human

Tissue Liver metastasis

Disease Adenocarcinoma

Metastatic site Liver

Characteristics

Age 77 years

Gender Male

Ethnicity Caucasian

Growth properties Adherent

HROC348Met Cells | 300871

Regulatory Data

Citation	HROC348Met (Cytion catalog number 300871)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_1U99

Biomolecular Data

MSI-status	MSS
-------------------	-----

Handling

Culture Medium	DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO ₃ (Cytion article number 820400a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Fluid renewal	Every 3 to 5 days
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HROC348Met Cells | 300871

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HROC348Met Cells | 300871

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.