

HROC103 T0 M1 Cells | 300802**General information**

Description	This is one of a series of cell lines which have been established by PD Dr. Michael Linnebacher from a PDTx (Patient-derived Tumor xenograft) since 2006.
Organism	Human
Tissue	Colorectal, Established from a PDx (patient-derived xenograft) of primary CRC tissue (Colon ascendens, TNM stage T2N1M0R0L0V0, grade G2, Lk(n) + 2, Σ Lk(n) 23).
Disease	Adenocarcinoma
Metastatic site	Regional lymph node involvement (TNM N1; Lk(n)+2 of 23 examined); no distant metastasis (M0)
Applications	Colorectal cancer research; CRC biology; PDx-derived cell line research; drug sensitivity and targeted therapy evaluation; p53/KRAS mutant CRC modeling; MSS CRC immunology; patient-matched HROC biobank studies
Synonyms	HROC103

Characteristics

Age	44 years
Gender	Male
Ethnicity	Caucasian
Morphology	Small cells in colonies
Cell type	Epithelial
Growth properties	Adherent

Regulatory Data

Citation	HROC103 T0 M1 (Cytion catalog number 300802)
Biosafety level	1
NCBI_TaxID	9606

HROC103 T0 M1 Cells | 300802**CellosaurusAccession** CVCL_1D10**GMO Status** No genetic modification; wildtype patient-derived CRC cell line established from a patient-derived xenograft by PD Dr. Linnebacher**Biomolecular Data****Ploidy status** Aneuploid**MSI-status** MSS**Mutational profile** P53 mut, APC mut, K-RasG12VA, N-Raswt, H-Raswt, PIK3CAwt, B-Rafwt**Handling****Culture Medium** DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)**Supplements** Supplement the medium with 10% FBS**Dissociation Reagent** Accutase**Doubling time** 30 hours**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Split ratio** 1 to 3**Seeding density** 2×10^4 cells/cm²**Fluid renewal** Every 3 to 5 days**Post-Thaw Recovery** Few days

HROC103 T0 M1 Cells | 300802

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

HROC103 T0 M1 Cells | 300802

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.