

## WT51 Cells | 302141

## General information

**Description** EBV-transformed B-lymphoblastoid cell line, derived from a male person, age unspecified. Homozygous cell line for HLA A:9, ,B:14, DR:4, and DP:2. Consanguineous parents. WT51 was part of the 10th International Histocompatibility Workshop (10IHW) cell line panel. Submitted by Dr.M.Trucco, HLA-Laboratory, Pittsburgh University Cancer Institute, USA.

**Organism** Human

**Tissue** Peripheral blood

**Applications** Functional analysis and genotyping of HLA Class II molecules. Analysis of B cell surface antigens, testing of cytotoxic drugs, mutational analysis, analysis of apoptotic mechanisms

**Synonyms** WT-51, WT 51, GM03103, GM3103, GM03103A

## Characteristics

**Age** Unspecified

**Gender** Male

**Ethnicity** Caucasian

**Morphology** Round cells

**Cell type** B lymphoblast

**Growth properties** Suspension

## Regulatory Data

**Citation** WT51 (Cytion catalog number 302141)

**Biosafety level** 1

**NCBI\_TaxID** 9606

**CellosaurusAccession** CVCL\_E887

**WT51 Cells | 302141**

**Biomolecular Data**

<b>Antigen expression</b>	CD19+
<b>Viruses</b>	Free of human pathogenic viruses SV40, JC/BK, HBV, HCV, and HIV. Contains EBV.
<b>Karyotype</b>	46, x,Y

**Handling**

<b>Culture Medium</b>	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO <sub>3</sub> (Cytion article number 820700a)
<b>Supplements</b>	Supplement the medium with 10% FBS
<b>Subculturing</b>	Gently homogenize the cell suspension in the flask by pipetting up and down, then take a representative sample to determine the cell density per ml. Dilute the suspension to achieve a cell concentration of $1 \times 10^5$ cells/ml with fresh culture medium, and aliquot the adjusted suspension into new flasks for further cultivation.
<b>Fluid renewal</b>	1 to 2 times per week
<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## WT51 Cells | 302141

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

**WT51 Cells | 302141**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.