

## L1210 Cells | 400257

### General information

#### Description

The L1210 cell line is a well-characterized murine lymphocytic leukemia model originally derived from a mouse with lymphoid leukemia. This cell line is widely used in cancer research due to its aggressive growth characteristics and high proliferative capacity. L1210 cells are commonly utilized in studies involving leukemia pathogenesis, chemotherapy drug testing, and the exploration of molecular mechanisms underlying cancer cell survival and proliferation.

L1210 cells exhibit rapid in vitro growth and maintain a suspension culture, making them ideal for in vitro assays and in vivo experiments, particularly in syngeneic mouse models. The cell line's responsiveness to a variety of chemotherapeutic agents has made it a valuable tool for preclinical screening of anti-leukemic drugs. Researchers often employ L1210 cells to study drug resistance mechanisms, evaluate novel therapeutic compounds, and investigate cellular responses to DNA-damaging agents.

Additionally, the L1210 cell line serves as a model to understand the immune response to leukemia, providing insights into how leukemia cells interact with the host's immune system. This includes studies on tumor immunology, cytokine production, and the efficacy of immunotherapeutic approaches. Overall, the L1210 cell line remains a critical resource in leukemia research, contributing to the advancement of cancer biology and therapeutic development.

**Organism** Mouse

**Tissue** Hematopoietic

**Disease** Leukemia

**Synonyms** L 1210, L-1210, Leukemic 1210, Leukemia 1210, Leukemia L1210

### Characteristics

**Breed/Subspecies** DBA/2

**Age** 8 months

**Gender** Female

**Cell type** Lymphoblast

**Growth properties** Suspension

### Regulatory Data

## L1210 Cells | 400257

<b>Citation</b>	L1210 (Cytion catalog number 400257)
-----------------	--------------------------------------

<b>Biosafety level</b>	1
------------------------	---

<b>NCBI_TaxID</b>	10090
-------------------	-------

<b>CellosaurusAccession</b>	CVCL_0382
-----------------------------	-----------

## Biomolecular Data

<b>Tumorigenic</b>	Yes, in nude mice and DBA mouse
--------------------	---------------------------------

<b>Viruses</b>	MAP-test negative: Sendai, Ektromelie, Polyoma, K-Virus, Kilham, Reo 3, PVM, LCM, M.pulmonis, MVM, Theiler's GD VII, Toolan's H-1, MHV, LDV, RCV/SDA, M-Adenovirus, B.piliformis.
----------------	---

## Handling

<b>Culture Medium</b>	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO <sub>3</sub> , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

<b>Supplements</b>	Supplement the medium with 10% horse serum
--------------------	--

<b>Doubling time</b>	10 to 12 hours
----------------------	----------------

<b>Subculturing</b>	Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of $5 \times 10^5$ cells/ml and keep the cell concentration within the range of $3 \times 10^5$ to $1 \times 10^6$ cells/ml for optimal growth.
---------------------	--

<b>Seeding density</b>	0.3 to $1 \times 10^6$ cells/ml
------------------------	---------------------------------

<b>Fluid renewal</b>	Every 3 to 4 days
----------------------	-------------------

<b>Post-Thaw Recovery</b>	Fast
---------------------------	------

<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	---

## L1210 Cells | 400257

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Flask Coating

None

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

**L1210 Cells | 400257**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.