

DSL-6B-C2 Cells | 500167

General information

Description

The DSL-6B/C2 cell line is derived from the DSL-6 transplantable acinar cell carcinoma of the pancreas, specifically established from a tumor model in a male Lewis rat. This model was initiated in 1986 from a primary acinar cell carcinoma that developed after intraperitoneal administration of azaserine, a potent carcinogen. The significance of this cell line stems from its origin in pancreatic cancer research, highlighting its utility in studying the biology and underlying mechanisms of pancreatic acinar cell carcinomas.

Initially, upon establishment in culture, DSL-6B/C2 cells exhibited the characteristic production of amylase, a hallmark of pancreatic exocrine function. However, this exocrine enzyme production was transient, ceasing within one to two weeks of culture. This change in phenotypic expression is notable as it suggests an adaptation to the in vitro environment, which may affect the cells' utility in certain types of biological assays. The loss of amylase production might also reflect changes in cell differentiation or the emergence of subpopulations within the cultured cells, which could be critical for researchers focusing on the evolution of tumor cell characteristics in vitro.

Organism Rat

Tissue Pancreas

Disease Carcinoma

Metastatic site Ductal

Synonyms DSL-6B/C2, DSL6B/C2

Characteristics

Breed/Subspecies Lewis

Age 2 years

Gender Male

Morphology Epithelial-like

Cell type Acinar cells

Growth properties Adherent

Regulatory Data

DSL-6B-C2 Cells | 500167

Citation DSL-6B-C2 (Cytion catalog number 500167)

Biosafety level 1

NCBI_TaxID 10116

CellosaurusAccession CVCL_4167

Biomolecular Data

Tumorigenic Yes, in Lewis rats the cells produce solid tumors and partially cystic tumors composed with a mixed phenotype of squamous, mucinous and glandular areas

Products Mucin

Handling

Culture Medium DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density 1×10^4 cells/cm² will yield in a confluent layer in about 4 days

Fluid renewal 2 times per week

Post-Thaw Recovery After thawing, plate the cells at 5×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

DSL-6B-C2 Cells | 500167

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Flask Coating

None

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

DSL-6B-C2 Cells | 500167

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.