

Human Dental Pulp Stem Cells (hDPSC) | 300702

General information

Description

Human Dental Pulp Stem Cells (DPSC, hDPSC) are multipotent stem cells isolated from the dental pulp of adult teeth, commonly third molars. These cells are particularly valuable in regenerative medicine due to their ability to differentiate into a variety of cell types, including those that form bone, cartilage, fat, and dental tissues. DPSCs are noted for their high proliferative capacity, making them a robust choice for tissue engineering and cell-based therapeutic applications.

DPSCs also possess significant immunomodulatory properties, which contribute to their potential use in treating inflammatory conditions. Beyond dental tissue regeneration, they have been investigated for their ability to repair bone defects and for their application in neurological therapies. Their relatively easy accessibility and the ability to maintain viability after cryopreservation make DPSCs an attractive option for clinical research and therapeutic development, particularly in the areas of regenerative dentistry, orthopedics, and neurodegenerative diseases.

Organism Human

Tissue Dental

Applications Drug testing, regenerative medicine, disease research

Characteristics

Growth properties Adherent

Regulatory Data

Citation Human Dental Pulp Stem Cells (DPSC, hDPSC) (Cytion catalog number 300702)

Biosafety level 1

NCBI_TaxID 9606

Biomolecular Data

Handling

Culture Medium Alpha MEM, w: 2.0 mM stable Glutamine, w/o: Ribonucleosides, w/o: Deoxyribonucleosides, w: 1.0 mM Sodium pyruvate, w: 2.2g/L NaHCO₃

Human Dental Pulp Stem Cells (hDPSC) | 300702

Supplements Supplement the medium with 10% FBS, 2 ng/mL bFGF

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Freeze medium As a cryopreservation medium, we use 90% FBS + 10% DMSO to maintain viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Human Dental Pulp Stem Cells (hDPSC) | 300702

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.