

T24 Cells | 300352**General information**

Description	Leukocytes and sera from patients with transitional cell carcinoma were cytotoxic to T24 and related lines.
Organism	Human
Tissue	Bladder
Disease	Carcinoma
Synonyms	T-24, T 24

Characteristics

Age	82 years
Gender	Female
Ethnicity	Caucasian
Morphology	Epithelial-like
Growth properties	Adherent

Regulatory Data

Citation	T24 (Cytion catalog number 300352)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0554

Biomolecular Data

Antigen expression	HLA A1, A3, B18, Bw35, Cw4, DRw2, Dw4
---------------------------	---------------------------------------

T24 Cells | 300352

Isoenzymes	Me-2, 1-2, PGM3, 1, PGM1, 1, ES-D, 1, AK-1, 1, GLO-1, 1, G6PD, B, Phenotype Frequency Product: 0.0216
Oncogenes	H-ras+
Tumorigenic	Yes, in hamster cheek pouch. Not in nude mice
Products	Tumor specific antigen
Karyotype	Hypodiploidy to hypopentaploidy, stemline 86, 2 to 4 telocentrics, 3 to 4 minutes, hypotetraploid to hypertetraploid with abnormalities including dicentrics, breaks, pulverization, minutes and telocentric markers
Handling	
Culture Medium	DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
Supplements	Supplement the medium with 5% FBS
Dissociation Reagent	Accutase
Doubling time	19 hours
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	1×10^4 cells/cm ²
Post-Thaw Recovery	After thawing, plate the cells at 5×10^4 cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

T24 Cells | 300352

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Flask Coating

None

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

T24 Cells | 300352

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.