

RAW 264.7 Cells | 400319

General information

Description

RAW 264.7 cells are a widely used murine macrophage cell line derived from the ascites of a male mouse with a tumor induced by the Abelson murine leukemia virus and are commonly used in immunological and infectious disease research. As an immortalized cell line, RAW264.7 cells are a key model system for studying macrophage biology, including immune responses to pathogens, signal transduction, and gene expression.

RAW264.7 cells are particularly valuable for their ability to differentiate into macrophage-like cells. These cells can be polarized into M1 macrophages, associated with inflammatory responses, or M2 macrophages, linked to tissue repair and anti-inflammatory processes. This polarization capacity, along with their ability to perform essential macrophage functions like pinocytosis and phagocytosis, underscores their relevance in studying macrophage biology and the complex interplay between immune responses and pathogens.

RAW 264.7 cells are instrumental in studying the immune system's interactions with various factors, including pathogens and bone biology. RAW264.7 cells can be induced to differentiate into osteoclast-like cells under certain conditions, such as exposure to RANKL (Receptor Activator of Nuclear Factor κ B Ligand), making them a model for studying certain aspects of osteoclast biology and bone resorption.

The RAW264.7 cell line's response to various stimuli, including the induction of pyroptosis, an inflammatory cell death process triggered by factors such as LPS (lipopolysaccharide), is instrumental in dissecting the pathways leading to inflammatory cytokine production. The impact of environmental conditions, such as extracellular glucose levels on cell function and phenotype, offers insights into cellular metabolism and the potential downregulation of inflammatory responses.

RAW264.7 cells, with their origins in murine leukemia and their extensive use in immunological research, serve as a crucial tool in advancing our understanding of macrophage biology, immune system-pathogen dynamics, osteoimmunology, and inflammatory responses, highlighting their indispensable role in both basic and applied biomedical research.

Organism

Mouse

Tissue

Ascites

Disease

Leukemia

Synonyms

RAW264, RAW2647, RAW264.7, RAW-264.7, Raw 264.7, Raw264.7

Characteristics

Age

Adult

Gender

Male

Cell type

Macrophage

RAW 264.7 Cells | 400319

Growth properties Adherent

Identifiers / Biosafety / Citation

Citation RAW 264.7 (Cytion catalog number 400319)

Biosafety level 2

Expression / Mutation

Receptors expressed Immunoglobulin (Fc), complement (C3)

Antigen expression H-2d

Viruses The cell line was tested and found positive for Reverse Transcriptase (RT) activity from C-Type retroviruses in cell culture supernatant and cell extract. Ectromelia virus (mousepox) may be secreted.

Products Lysozyme

Handling

Culture Medium RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Medium supplements Supplement the medium with 10% FBS

Passaging solution Accutase

Doubling time RAW264.7 cells exhibit a doubling time ranging from 11 to 30 hours

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

RAW 264.7 Cells | 400319

Seeding density 4 x 10⁴ cells/cm²

Fluid renewal 2 to 3 times per week

Freeze medium CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

RAW 264.7 Cells | 400319

STR profile

Amelogenin: x,y

M_18-3: 18

M_4-2: 22.3,23.3

M_6-7: 12

M_3-2: 14

M_19-2: 12,14

M_7-1: 25.2

M_1-1: 15,16

M_8-1: 13

M_2-1: 16

M_15-3: 22.3

M_6-4: 18

M_11-2: 17

M_1-2: 17

M_17-2: 14,16

M_12-1: 16,17

M_5-5: 14

M_X-1: 25

M_13-1: 16.2

Human D4/D8: -