

A498 Cells | 300113

General information

Description

A498 cells are a human renal cell carcinoma cell line derived from the kidney tissue of a 58-year-old Caucasian male. These cells are extensively used in research related to kidney cancer, particularly for studying clear cell renal cell carcinoma, which is the most common type of kidney cancer in adults.

The A498 cell line is characterized by its epithelial-like morphology and has been a valuable model for investigating the molecular and cellular mechanisms of renal carcinogenesis. These cells exhibit several features typical of kidney cancer, including alterations in the expression of genes involved in cell cycle regulation, apoptosis, and angiogenesis.

A498 cells are particularly useful for examining the metabolic pathways altered in kidney cancer, as they display a distinct metabolic profile that includes changes in lipid and glucose metabolism. This aspect makes them suitable for metabolic targeting studies, which explore how altering metabolic pathways can inhibit tumor growth.

Furthermore, A498 cells are employed in drug discovery and toxicology studies to test the efficacy of new chemotherapeutic agents and targeted therapies. They are also used to study the response of renal cancer cells to hypoxic conditions, a common feature of solid tumors that significantly influences tumor behavior and treatment response.

Overall, the A498 cell line serves as an essential tool in renal cancer research, facilitating the development of more effective therapeutic strategies and enhancing our understanding of kidney cancer biology.

Organism Human

Tissue Kidney

Disease Renal cell carcinoma

Synonyms A-498

Characteristics

Age 52 years

Gender Male

Ethnicity Caucasian

Morphology Epithelial-like

Growth properties Monolayer, adherent

A498 Cells | 300113

Identifiers / Biosafety / Citation

Citation	A498 (Cytion catalog number 300113)
Biosafety level	1

Expression / Mutation

Isoenzymes	PGM3, 1, PGM1, 1-2, ES-D, 2, Me-2, 1, AK-1, 1, GLO-1, 2, G6PD, B
Tumorigenic	Yes, in nude mice. Forms undifferentiated carcinoma, also forms tumors in anti thymocyte serum treated newborn mice
Ploidy status	Bimodal, tetraploid

Handling

Culture Medium	EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Doubling time	62 hours
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Split ratio	A ratio of 1:2 to 1:4 is recommended
Seeding density	1 x 10 ⁴ cells/cm ² will result in a confluent monolayer within 4 days.
Fluid renewal	Every 3 days

A498 Cells | 300113

Freezing recovery

After thawing, plate the cells at 2×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 to 48 hours.

Freeze medium

CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

A498 Cells | 300113

STR profile

Amelogenin: x,x
CSF1PO: 11,12
D13S317: 12
D16S539: 12
D5S818: 11,13
D7S820: 11,12
TH01: 6,9,3
TPOX: 8,11
vWA: 18
D3S1358: 15
D21S11: 28,32
D18S51: 17
Penta E: 10,14
Penta D: 9,14
D8S1179: 13,15
FGA: 18,2

HLA alleles

A*: 02:01:01
B*: 08:01:01
C*: 07:01:01
DRB1*: 03:01:01
DQA1*: 05:01:01
DQB1*: 02:01:01
DPB1*: 01:01:01
E: 01:03:02