

LNCaP Cells | 300265

General information

Description

LNCaP cells, derived from a metastatic lesion in a lymph node of a prostate cancer patient, represent a critical tool in prostate cancer research, particularly for studying the role of androgens and androgen receptor (AR) dynamics in cancer progression. The LNCaP cell line is characterized by their androgen-sensitive growth and offers a window into the mechanisms underlying prostate cancer's response to hormonal manipulation.

As a model for metastatic prostate cancer, parental LNCaP cells and their sublines, such as the LNCaP clone FGC, provide clinically relevant insights into disease progression, especially in the context of metastasis to bone, forming osteoblastic lesions akin to those observed in human prostate cancer.

The LNCaP human prostate cancer cell line expresses a mutated form of the AR gene with broader steroid-binding specificity and therefore is pivotal for understanding the complex interplay between AR activity and prostate cancer progression. This includes the examination of AR downstream targets like PSA and NKx3.1, which are crucial for prostatic epithelial cell function. LNCaP cells are further used in cytotoxicity studies such as those induced by ripl or the potential therapeutic effects of compounds like amygdalin, within the scope of intracellular drug delivery strategies.

In summary, the human prostate carcinoma cell line LNCaP serves as a cornerstone in understanding the role of androgens in cancer progression and prostate cancer, offering insights into hormone-responsive cancers, the challenges of resistant prostate cancer, and the potential for therapeutic interventions. The LNCaP cell line is considered one of the classic and most widely used human prostate cancer cell lines, alongside DU145 and PC3 cells.

Organism	Human
Tissue	Prostate
Disease	Carcinoma
Metastatic site	Left supraclavicular lymph node
Synonyms	LNCAP, LNCap, Ln-Cap, Lymph Node Carcinoma of the prostate

Characteristics

Age	50 years
Gender	Male
Ethnicity	Caucasian
Morphology	Epithelial-like

LNCaP Cells | 300265

Growth properties Adherent, clusters

Identifiers / Biosafety / Citation

Citation LNCaP (Cytion catalog number 300265)

Biosafety level 1

Expression / Mutation

Receptors expressed Androgen, estrogen

Protein expression p53 positive

Tumorigenic Yes, in nude mice

Products human prostatic acid phosphatase, prostate specific antigen

Karyotype pseudodiploid male, seven marker chromosomes, modal number = 46, range = 33 to 91

Handling

Culture Medium EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO₃, w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)

Medium supplements Supplement the medium with 10% FBS

Passaging solution Accutase

Doubling time 60 hours

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

LNCaP Cells | 300265

Split ratio A ratio of 1:3 to 1:6 is recommended

Seeding density 1 to 2 x 10⁴ cells/cm²

Fluid renewal Every 3 days

Freezing recovery After thawing, plate the cells at 5 x 10⁴ cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

LNCaP Cells | 300265

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

Amelogenin: x,y
CSF1PO: 10,11
D13S317: 10,12
D16S539: 11
D5S818: 11,12
D7S820: 9.1,10.3
TH01: 9
TPOX: 8,9
vWA: 16,18
D3S1358: 16
D21S11: 29,31.2
D18S51: 11,12
Penta E: 12,16
Penta D: 12,12.4
D8S1179: 12,14
FGA: 19,2

HLA alleles

A*: 01:01:01, 02:01:01
B*: 08:01:01, 37:01:01
C*: 06:02:01, 07:01:01
DRB1*: 03:01:01, 10:01:01
DQA1*: 01:05:01, 05:01:01
DQB1*: 02:01:01, 05:01:01
DPB1*: 02:01:02G, 04:02:01G
E: 01:01:01