

A704 Cells | 300217**General information****Description**

A-704 is a human epithelial cell line derived from kidney tissue of a 78-year-old male patient with adenocarcinoma. This cell line exhibits an epithelial morphology. It is a valuable resource in cancer research, particularly for studying adenocarcinoma. A-704 is a versatile cell line with applications in 3D cell culture and as a transfection host.

Derived by D.J. Giard, A-704 maintains consistency and reliability in experimental settings. Karyotype analysis reveals that A-704 cells exhibit abnormalities such as breaks, dicentrics, and endoreduplication, ranging from diploid to hyperdiploid, hypertriploid to hypertetraploid.

While not tumorigenic in immunosuppressed mice, A-704 cells can form colonies in a semisolid medium. A-704 cells exhibit specific isoenzyme profiles, including AK-1, ES-D, G6PD, GLO-I, Me-2, PGM1, and PGM3.

Organism Human**Tissue** Kidney**Disease** Adenocarcinoma**Synonyms** A.704, A-704**Characteristics****Age** 78 years**Gender** Male**Ethnicity** Caucasian**Morphology** Epithelial-like**Growth properties** Monolayer, adherent**Identifiers / Biosafety / Citation****Citation** A704 (Cytion catalog number 300217)**Biosafety level** 1

A704 Cells | 300217

Expression / Mutation

Isoenzymes	Me-2, 1, PGM3, 1-2, PGM1, 1, ES-D, 1, AK-1, 1, GLO-1, 2, G6PD, B
Tumorigenic	No
Karyotype	(P59) diploid to hyperdiploid, hypertriploid to hypertetraploid with abnormalities including breaks, dicentrics and endoreduplication

Handling

Culture Medium	EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Split ratio	A ratio of 1:3 to 1:4 is recommended
Seeding density	1 x 10 ⁴ cells/cm ² will result in a confluent monolayer within 4 days.
Fluid renewal	2 to 3 times per week
Freezing recovery	After thawing, plate the cells at 5 x 10 ⁴ cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

A704 Cells | 300217

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

A704 Cells | 300217

STR profile

Amelogenin: x,y
CSF1PO: 7,8
D13S317: 8
D16S539: 12,13
D5S818: 10,11
D7S820: 10
TH01: 7,9
TPOX: 11
vWA: 14,18
D3S1358: 15
D21S11: 28,32
D18S51: 16,17
Penta E: 8,17
Penta D: 2,2,11
D8S1179: 13,15
FGA: 22,23

HLA alleles

A*: 34:02:01, 74:01:01
B*: 35:01:01, 44:03:01
C*: 04:01:01
DRB1*: 15:03:01G
DQA1*: 01:02:01
DQB1*: 06:02:01
DPB1*: 02:01:19, 04:02:01G
E: 01:01:01, 01:03