

U2OS Cells | 300364

General information

Description

U2OS cells, an osteosarcoma cell line derived from a human osteosarcoma patient, play a significant role in cancer research, particularly in the study of bone cancer. U2OS cells are used extensively in cancer research, drug development, apoptosis studies, genetic research, and radiation oncology studies. The value of U2OS cells lies in their application to investigate apoptosis and drug resistance, essential for creating small molecule inhibitors and similar therapeutic agents.

In the realm of clinical osteosarcoma research, the U2OS cell line is instrumental in examining biological responses to radiotherapy, thereby enriching our understanding of osteosarcoma biology. These cells are also pivotal in investigating chromatin modifications and their impact on cell biology, especially in the context of tumor formation and cancer progression.

The U2OS cell line, also referred to as the OS cell line, is recognized for its *in vivo* tumor formation capacity when administered through subcutaneous and intramuscular injections. The tumors produced by U2OS cells are characterized as high-grade sarcomas and exhibit significant osteoid production, which is a hallmark of osteosarcoma. Additionally, these tumors showed infiltration by immune cells. U2OS therefore serves as a representative model for studying human osteosarcoma, its interactions with the human immune system and tumor immunology. One of the challenges, however, is ensuring the osteosarcoma U2OS cell line accurately reflects the tumors *in vivo*, given the variability in tumor formation capacity.

In summary, sarcoma cell lines such as U2OS serve as a pivotal tool in understanding osteosarcoma, offering valuable insights into cancer biology, therapeutic development, and the complexities of tumor-immune system interactions, while highlighting the need for accurate *in vivo* tumor modeling.

Organism Human

Tissue Bone, tibia

Disease Osteosarcoma

Synonyms U-2 OS, U-2OS, U-2-OS, U2-OS, U20-S, U20S, 2T

Characteristics

Age 15 years

Gender Female

Ethnicity Caucasian

Morphology Epithelial-like

U2OS Cells | 300364

Growth properties	Monolayer, adherent
--------------------------	---------------------

Identifiers / Biosafety / Citation

Citation	U-2 OS (Cytion catalog number 300364)
-----------------	---------------------------------------

Biosafety level	1
------------------------	---

Depositor	Lee
------------------	-----

Expression / Mutation

Receptors expressed	Insulin-like growth factor I (IGF-I), insulin-like growth factor II (IGF-II), Osteosarcoma derived growth factor (ODGF)
----------------------------	---

Antigen expression	Blood Type A, Rh+, HLA A2, Aw30, B12, Bw35, B40(+/-)
---------------------------	--

Isoenzymes	PGM3, 1, PGM1, 2, ES-D, 1, AK-1, 1, GLO-1, 2, G6PD, B, Phenotype Frequency Product: 0.0082
-------------------	--

Products	Osteosarcoma derived growth factor (ODGF)
-----------------	---

Karyotype	(P11-46) hypodiploid to near tetraploid, (P111-118) modal numbers 34 to 37 and 64 to 67 with abnormalities including dicentrics, breaks, rings, and pulverizations plus acrocentric subtelocentric and minute markers
------------------	---

Handling

Culture Medium	DMEM:Ham's F12, w: 3.1 g/L Glucose, w: 1.6 mM L-Glutamine, w: 15 mM HEPES, w: 1.0 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
-----------------------	---

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Passaging solution	Accutase
---------------------------	----------

U2OS Cells | 300364

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Split ratio A ratio of 1:3 to 1:6 is recommended

Seeding density 1×10^4 cells/cm²

Fluid renewal 2 to 3 times per week

Freeze medium CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

U2OS Cells | 300364

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

CSF1PO: 12,13

D13S317: 13

D16S539: 11,12

D5S818: 8,11

D7S820: 11,12

TH01: 6,9,3

TPOX: 11,12

vWA: 14,18

D3S1358: 16

D21S11: 31

D18S51: 12,14

D8S1179: 12,14

FGA: 20

D2S1338: 20,24

D19S433: 15

HLA alleles

A*: 02:01:01, 32:01:01

B*: 44:02:01, 44:27:01

C*: 05:01:01, 07:04:01

DRB1*: 09:01:02, 14:54:01

DQA1*: 01:04:01, 03:02:01

DQB1*: 03:03:02, 05:03:01

DPB1*: 02:01:02, 04:01:01

E: 01:01:01