

HROC103 T0 M1 Cells | 300802

General information

Description	This is one of a series of cell lines which have been established by PD Dr. Michael Linnebacher from a PDTx (Patient-derived Tumor xenograft) since 2006.
Organism	Human
Tissue	Colorectal, Established from a PDx (patient-derived xenograft) of primary CRC tissue (Colon ascendens, TNM stage T2N1M0R0L0V0, grade G2, Lk(n) + 2, ? Lk(n) 23).
Disease	Adenocarcinoma
Synonyms	HROC103

Characteristics

Age	44 years
Gender	Male
Ethnicity	Caucasian
Morphology	Small cells in colonies
Cell type	Epithelial
Growth properties	Adherent

Identifiers / Biosafety / Citation

Citation	HROC103 T0 M1 (Cyton catalog number 300802)
Biosafety level	1
Depositor	M. Linnebacher

Expression / Mutation

Ploidy status	Aneuploid
----------------------	-----------

HROC103 T0 M1 Cells | 300802

Mutational profile p53 mut, APC mut, K-RasG12VA, N-Raswt, H-Raswt, PIK3CAwt, B-Rafwt

Handling

Culture Medium	DMEM:Ham's F12, w: 3.1 g/L Glucose, w: 1.6 mM L-Glutamine, w: 15 mM HEPES, w: 1.0 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Doubling time	30 hours
Subculturing	Remove medium and rinse the adherent cells using PBS without calcium and magnesium (3-5 ml PBS for T25, 5-10ml for T75 cell culture flasks). Add TrypLE Express (1-2ml per T25, 2.5ml per T75 cell culture flask), the cell sheet must be covered completely. Incubate at 37 degree Celsius for 10 to 15 minutes. Carefully resuspend the cells with medium (10 ml), centrifuge for 3 min at 300xg, resuspend cells in fresh medium and dispense into new flasks which contain fresh medium.
Split ratio	A ratio of 1:3 to 1:5 is recommended
Seeding density	2 x 10 ⁴ cells/cm ²
Fluid renewal	Every 3 to 5 days
Freezing recovery	Few days
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

HROC103 T0 M1 Cells | 300802

Handling of cryopreserved cultures

HROC103 T0 M1 cells are shipped in a deep-frozen state on dry ice. Upon receipt, confirm that the vial remains frozen. For storage, place the cryovial immediately at temperatures below -150 degrees. If you plan to culture the cells immediately, swiftly thaw the vial by shaking it in a 37 degrees water bath with clean water and an antimicrobial agent for 40-60 seconds. Remove the vial once a small ice clump persists, ensuring it remains cold. Proceed with all subsequent steps under aseptic conditions. In a sterile flow hood, disinfect the cryovial with 70% ethanol. Then, gently open the vial and transfer the cell suspension into a 15 ml centrifuge tube pre-filled with 8 ml of room temperature culture medium. Gently mix the cells. For cell separation, centrifuge at 300 x g for 3 minutes and dispose of the supernatant. Skipping centrifugation is optional, although any residual freezing medium should be removed after 24 hours. Resuspend the pellet gently in 10 ml of fresh culture medium and divide between two T25 culture flasks. Follow the subculture protocol for subsequent steps.

Handling of proliferating cultures

One or two cell culture flasks come filled with cell culture medium. Collect the entire medium in a 50 ml centrifuge tube. Spin down the collected medium at 300 x g for 3 minutes to collect the cells which may have detached during transit. If a cell pellet is visible, resuspend the cells in 5 ml of cell culture medium and transfer to a T25 cell culture flask. Carefully add 5 ml of cell culture medium to each T25 cell culture flask. Examine cell morphology and confluency using a microscope. Finally, incubate the flasks at 37 degrees Celsius for at least 24 hours.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is rigorously excluded using both PCR-based assays and luminescence-based mycoplasma detection methods. To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

Amelogenin: x (Patient male, Y lost)
CSF1PO: 12
D13S317: 11,12
D16S539: 12,13
D5S818: 12,13
D7S820: 9
TH01: 6,9
TPOX: 8
vWA: 14,15