

T84 Cells | 300354

General information

Description	This line exhibits tight junctions, and desmosomes between adjacent cells. The cells should be maintained at high density (at least 1/4 confluency).
Organism	Human
Tissue	Colon
Disease	Carcinoma
Metastatic site	Lung
Synonyms	T-84, T 84

Characteristics

Age	72 years
Gender	Male
Morphology	Epithelial-like
Growth properties	Adherent

Identifiers / Biosafety / Citation

Citation	T84 (Cytion catalog number 300354)
Biosafety level	1

Expression / Mutation

Receptors expressed	Peptide hormone, neurotransmitter
Antigen expression	Keratin + (Immunoperoxidase staining)
Isoenzymes	G6PD, B, PGM1, 1, PGM3, 1, ES-D, 1, Me-2, 1-2, AK-1, 1, GLO-1, 1-2

T84 Cells | 300354

Tumorigenic	Yes, in nude mice
Products	Carcinoembryonic antigen (CEA), 600 ng/ml per 10 exp6 cells per 10 days, keratin
Mutational profile	T84 cells carry a heterozygous Kras mutation in codon13: GGC(Wt Gly) >GAC(Asp)
Karyotype	The stemline modal chromosome number is 56, occurring at 28% with polyploidy at 12.4%. Eighteen markers are common to most metaphases examined. Normal x and chromosome 13 were absent, chromosomes 2, 4 and 22 were single-copied, and chromosome 12 was 4-copied.No Y chromosome was detected by Q band observation. DM occurred in nearly 50% of the cells.

Handling

Culture Medium	Ham's F12, w: 1.0 mM stable Glutamine, w: 1.0 mM Sodium pyruvate, w: 1.1 g/L NaHCO3 (Cytion article number 820600a)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Split ratio	A ratio of 1:2 to 1:4 is recommended
Fluid renewal	2 times per week
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

T84 Cells | 300354

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

T84 Cells | 300354

STR profile

CSF1PO: 10
D13S317: 9
D16S539: 10,11
D5S818: 12
D7S820: 8,10
TH01: 6,9
TPOX: 8
vWA: 17,18
D3S1358: 19
D21S11: 31
D18S51: 17
Penta E: 14
Penta D: 9
D8S1179: 15
FGA: 24

HLA alleles

A*: 02:01:01, 24:02:01
B*: 18:01:01, 35:01:01
C*: 04:01:01, 07:01:01
DRB1*: 01:01:01, 09:01:02
DQA1*: 01:01:01, 03:02:01
DQB1*: 03:03:02, 05:01:01
DPB1*: 02:01:02, 04:01:01
E: 01:03:01, 01:03:02