

## P-815 Cells | 400242

### General information

**Description** P-815 cells phagocytose latex beads but not zymosan or BCG. They do not function in antibody dependent cell mediated cytotoxicity. Growth of the cells is not inhibited by dextran sulfate, LPS or PPD. Tested and found negative for ectromelia virus (mousepox).

**Organism** Mouse

**Tissue** Hematopoietic

**Disease** Mastocytoma

**Synonyms** P815, P 815

### Characteristics

**Gender** Male

**Morphology** Round cells

**Cell type** Mast cell

**Growth properties** Suspension, some adherent cells

### Identifiers / Biosafety / Citation

**Citation** P-815 (Cytion catalog number 400242)

**Biosafety level** 1

**Depositor** Ralph

### Expression / Mutation

**Products** Lysozyme

### Handling

### P-815 Cells | 400242

<b>Culture Medium</b>	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO <sub>3</sub> , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

<b>Medium supplements</b>	Supplement the medium with 10% FBS
---------------------------	------------------------------------

<b>Subculturing</b>	Start cultures at $2 \times 10^5$ cells/ml and maintain between $1 \times 10^5$ and $1 \times 10^6$ cells/ml. Subculture by transferring an appropriate aliquot of the suspension into new flasks filled with fresh cell culture medium. Detach adherent cells by washing with PBS first and incubation with Accutase at ambient temperatur 8-10 minutes. Combine with the nonadherent cells and distribute into new cell culture flasks.
---------------------	---

<b>Seeding density</b>	1 to $2 \times 10^6$ cells/ml
------------------------	-------------------------------

<b>Fluid renewal</b>	2 to 3 times per week
----------------------	-----------------------

<b>Freezing recovery</b>	After thawing, allow the cells to recover from the freezing process for at least 24 hours.
--------------------------	--

<b>Freeze medium</b>	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
----------------------	--

### P-815 Cells | 400242

#### Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Quality control / Genetic profile / HLA

#### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

#### STR profile

**Amelogenin:** x,x