

EMT6 Cells | 305159

General information

Description

The EMT6 cell line is a murine mammary adenocarcinoma line that is extensively used in cancer research, particularly in studies related to breast cancer. Originating from a spontaneous tumor in a BALB/c mouse, EMT6 cells are employed both in vitro and in vivo to analyze tumorigenesis, metastasis, and chemotherapeutic resistance. The cells are characterized by their ability to form tumors rapidly when transplanted into immunocompetent mice, making them an ideal model for studying tumor immunity and the efficacy of anti-cancer therapies.

EMT6 cells are highly adaptable to various growth conditions and have a relatively high mitotic index, which facilitates easy cultivation and experimental manipulation in laboratory settings. They are also used in radiobiology studies due to their pronounced sensitivity to radiation, providing insights into the cellular mechanisms underlying radiation therapy for cancer. The cell line has been instrumental in the development of protocols for hypoxic cell sensitizers and has been used to test the efficacy of photodynamic therapy agents.

Organism

Mouse

Tissue

Breast

Disease

Malignant neoplasms of the mouse mammary gland

Synonyms

EMT-6, Experimental Mammary Tumour-6

Characteristics

Breed/Subspecies

BALB/cCRGL

Gender

Female

Morphology

Epithelial

Growth properties

Adherent

Regulatory Data

Citation

EMT6 (Cytion catalog number 305159)

Biosafety level

1

NCBI_TaxID

10090

EMT6 Cells | 305159

CellosaurusAccession CVCL_1923

Biomolecular Data**Handling**

Culture Medium DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Fluid renewal 2 to 3 times per week

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

EMT6 Cells | 305159

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

EMT6 Cells | 305159

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.