LLC-PK1 Cells | 607264



General information

Description	 LLC-PK1 cells are a well-established and widely used cell line in biomedical research. These cells were derived from a healthy male pig's kidney, exhibiting typical epithelial morphology. The LLC-PK1 line is polarized and contains tight junctions, making it an ideal model for epithelial tissue. One of the critical features of LLC-PK1 cells is their ability to produce plasminogen activator, a substance that stimulates fibrinolysis. This property has made LLC-PK1 cells particularly valuable in thrombosis research. In recent years, plasminogen activator has been included in drugs used in thrombosis therapies since it facilitates the dissolution of small blood clots. In addition to producing plasminogen activators, LLC-PK1 cells produce large amounts of cytokeratin. This characteristic has made them popular for various pharmacologic and metabolic research investigations. The LLC-PK1 line has been used in drug metabolism, transport, toxicity, and interaction studies. LLC-PK1 cells are also frequently used in permeability assays. The mechanism of uracil transport differs depending on cell lines, with a Na+-independent system on the basolateral membrane in Caco-2 cells and both Na+-dependent and Na+-independent systems on the apical membrane in LLC-PK1 cells. Compared to other cell lines, LLC-PK1 cells share many characteristics of proximal tubular cells in vivo, including apical membrane incrovilli, high activities of apical membrane enzymes, and expression of parathyroid hormone receptors and sodium-dependent glucose transporters. This makes LLC-PK1 cells avaluable tool in renal toxicology studies. Another cell line commonly used in renal toxicology studies is the MDCK cell line. Like LLC-PK1 cells are also capable of forming 'domes', fluid-filled blisters resulting from water and solute transport, tight junctions, and adhesion of the cells to the substratum. In conclusion, the LLC-PK1 cells are also capable of forming 'domes', fluid-filled blisters resulting fro
Organism	Sus Scrofa
Tissue	Kidney
Applications	Drug metabolism, permeability assays, toxicity, and interaction studies.
Synonyms	LLC-PK(1), LLC-PK-1, LLC PK-1, LLC-PK1, LLC PK1, LLCPK1, Lilly Laboratories Cell-Porcine Kidney 1

Characteristics

Age

3-4 weeks



LLC-PK1 Cells | 607264

Gender	Male
Morphology	Epithelial-like
Growth properties	Adherent/suspension. It takes a couple of days until cells grow in adherent colonies.

Identifiers / Biosafety / Citation

Citation	LLC-PK1 (Cytion catalog number 607264)
Biosafety level	1

Expression / Mutation

Products	plasminogen activator

Handling

Culture Medium	DMEM:Ham's F12, w: 3.1 g/L Glucose, w: 1.6 mM L-Glutamine, w: 15 mM HEPES, w: 1.0 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
Medium supplements	Supplement the medium with 10% FBS
Passaging solution	Accutase
Subculturing	Gather the suspension cells in a 15 ml tube and gently wash the adherent cells with PBS lacking calcium and magnesium (use 3-5 ml for T25 flasks and 5-10 ml for T75 flasks). Apply Accutase (1-2 ml for T25 flasks, 2.5 ml for T75 flasks) ensuring full coverage of the cell layer. Allow the cells to incubate at room temperature for 10 minutes. Following incubation, combine and centrifuge both the suspension and adherent cells. After centrifugation, carefully resuspend the cell pellet and transfer the cell suspension into new flasks containing fresh medium.
Split ratio	A ratio of 1:3 to 1:8 is recommended
Seeding density	1 to 3 x 10^6 cells/cm^2
Fluid renewal	Every 3 days

Product sheet

LLC-PK1 Cells | 607264



Freezing recovery	After thawing, plate the cells at 5 x 10^4 cells/cm^2 and allow the cells to recover from the freezing process and to adhere for at least 24 hours.
Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
Handling of cryopreserved cultures	1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
	2. Upon receipt, either store the cryovial immediately at temperatures below -150?C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
	3. For immediate culturing, swiftly thaw the vial by immersing it in a 37?C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
	4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
	5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
	6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
	7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
	8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.