

HuCC-T1 Cells | 300469

General information

Organism	Human
Tissue	Liver
Disease	Intrahepatic cholangiocarcinoma
Metastatic site	Ascites
Applications	Studies of the mechanism of tumor marker secretion and tumor cell growth in the human cholangiocellular carcinoma
Synonyms	HuCC-T1, HUCCT-1, HUCC-T1, HUCCT1, HuCCT1

Characteristics

Age	56 years
Gender	Male
Ethnicity	Japanese
Morphology	Epithelial
Growth properties	Adherent

Identifiers / Biosafety / Citation

Citation	HuCC-T1 (Cytion catalog number 300469)
Biosafety level	1

Expression / Mutation

Tumorigenic	Yes, in nude mice.
--------------------	--------------------

Handling

HuCC-T1 Cells | 300469

Culture Medium RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Medium supplements Supplement the medium with 10% FBS

Passaging solution Trypsin-EDTA

Subculturing Discard the old medium and wash the cells with PBS. Add a freshly prepared 0.025% trypsin/0.02% EDTA solution heated to 37 degrees Celsius and wait until the cells detach, which usually takes about 5 minutes. Neutralize the trypsin by adding fresh medium, then transfer the cell mixture to a tube and centrifuge. After centrifugation, remove the supernatant, resuspend the cell pellet in fresh culture medium, and transfer the suspension to new flasks. Incorporate G418 into the culture medium to achieve a final concentration of 0.5 mg/ml

Freeze medium CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

HuCC-T1 Cells | 300469

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

HuCC-T1 Cells | 300469

STR profile

Amelogenin: x,y
CSF1PO: 11,12
D13S317: 11,13
D16S539: 11,12
D5S818: 12,13
D7S820: 10,11
TH01: 7,10
TPOX: 8
vWA: 18
D3S1358: 15
D21S11: 31
D18S51: 13
Penta E: 15,18
Penta D: 10
D8S1179: 10
FGA: 20,23
D6S1043: 13
D2S1338: 17,18
D12S391: 18,20
D19S433: 13