

Jiyoye Cells | 300366**General information**

Description	Established from a 7-year-old boy with Burkitt lymphoma. Epstein-Barr virus positive but according to TRBA 468 stable in the latent phase. The subline P3HR-1, deficient in HLA class II, is also available at CLS (order number 302016).
Organism	Human
Tissue	Lymphatic System
Disease	B-cell Non-Hodgkin-Lymphoma
Metastatic site	B-Lymphocyte
Applications	Analysis of B cell surface antigens, testing of cytotoxic drugs, mutational analysis, analysis of apoptotic mechanisms, haplotype standard.
Synonyms	JIYOYE, Jijoye, JIJOYE, P-2003, P3 (Jiyoye), P-3-Jijoye, P3-Jiyoye, P-3J, P3J, Jiyoye(P-2003), Jiyoye (P-2003), JiyoyeP-2003, OB2, GM04678

Characteristics

Age	7 years
Gender	Male
Ethnicity	African
Cell type	B lymphocyte
Growth properties	Suspension

Identifiers / Biosafety / Citation

Citation	Jiyoye (Cytion catalog number 300366)
Biosafety level	1

Expression / Mutation

Jiyoye Cells | 300366

Antigen expression	CD10+, CD19+
---------------------------	--------------

Karyotype	46, hypodiploid
------------------	-----------------

Handling

Culture Medium	RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Medium supplements	Supplement the medium with 10% FBS
---------------------------	------------------------------------

Subculturing	Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 2×10^5 cells/ml and keep the cell concentration within the range of 1×10^5 to 1×10^6 cells/ml for optimal growth.
---------------------	--

Seeding density	3×10^5 cells/ml
------------------------	--------------------------

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freezing recovery	Fast (48 hours)
--------------------------	-----------------

Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
----------------------	--

Jiyoye Cells | 300366

Handling of cryopreserved cultures

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

Jiyoye Cells | 300366

STR profile

Amelogenin: x,y
CSF1PO: 10,11
D13S317: 12
D16S539: 10,11
D5S818: 12
D7S820: 8,10
TH01: 7,9
TPOX: 6,8
vWA: 15,19
D3S1358: 16,17
D21S11: 28,36
D18S51: 12
Penta E: 8,12
Penta D: 2.2,12
D8S1179: 14,15
FGA: 23,24

HLA alleles

A*: 03:01:01, 03.01.1900 02:01
B*: 02.01.1900 05:01, 02.01.1900 10:01
C*: 04:01:01
DRB1*: 11:02:01, 15:03:01
DQA1*: 01:02:01, 05:05:01
DQB1*: 03:19:01, 06:02:01
DPB1*: 01:01:01, 02:01:02
E: 01:01, 01:03