

HT-1080 Cells | 300216

General information

Description

HT-1080 cells, derived from the connective tissue of a 35-year-old male patient with Fibrosarcoma in 1972, are widely used for studying the mechanisms of tumor invasiveness and metastasis due to their highly aggressive and invasive nature.

HT-1080 cells have been extensively utilized in studies involving cell migration, invasion assays, and the testing of anti-cancer compounds. In the realm of therapeutic development, HT-1080 cells are employed in the screening of anti-cancer drugs and in the evaluation of their effects on cell viability, apoptosis, and metastatic potential.

HT-1080 cells have also been used in research focusing on the extracellular matrix, angiogenesis, and the role of various genes and proteins in cancer progression. HT-1080 cells produce matrix metalloproteinases (MMPs), enzymes that degrade components of the extracellular matrix and play a critical role in tumor invasion and metastasis. This feature makes the HT-1080 cell line useful for studies investigating the regulation of MMPs and their inhibitors.

In summary, the HT-1080 cell line, with its extensive applications in the study of cancer research, cell adhesion, migration, and invasion models, as well as in the development of therapeutic strategies, continues to be a valuable resource in cancer research.

Organism Human

Disease Fibrosarcoma

Synonyms Ht-1080, HT 1080, HT1080, HT 1080.T

Characteristics

Age 35 years

Gender Male

Ethnicity Caucasian

Morphology Epithelial-like

Cell type Fibroblast

Growth properties Adherent

Identifiers / Biosafety / Citation

HT-1080 Cells | 300216

Citation	HT-1080 (Cytion catalog number 300216)
-----------------	----------------------------------------

Biosafety level 1

Expression / Mutation

Isoenzymes	G6PD, B
-------------------	---------

Oncogenes ras+

Tumorigenic	Yes, in immunosuppressed mice
--------------------	-------------------------------

Virus susceptibility poliovirus 1, vesicular stomatitis (Indiana), RD114, feline leukemia virus (FeLV)

Reverse transcriptase	negative
------------------------------	----------

Karyotype Modal number: 2n=46, pseudodiploid

Handling

Culture Medium	EMEM, w: 2 mM L-Glutamine, w: 1.5 g/L NaHCO ₃ , w: EBSS, w: 1 mM Sodium pyruvate, w: NEAA (Cytion article number 820100c)
-----------------------	--------------------------------------------------------------------------------------------------------------------------------------

Medium supplements Supplement the medium with 10% FBS

Passaging solution	Accutase
---------------------------	----------

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Split ratio	A ratio of 1:4 to 1:8 is recommended
--------------------	--------------------------------------

Seeding density 1 x 10⁴ cells/cm²

HT-1080 Cells | 300216

Fluid renewal	Every 3 days
Freezing recovery	After thawing, plate the cells at 5×10^4 cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium	CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)
----------------------	------------------------------------------------------------------------------

Handling of cryopreserved cultures

HT-1080 cells are shipped in a deep-frozen state on dry ice. Upon receipt, confirm that the vial remains frozen. For storage, place the cryovial immediately at temperatures below -150 degrees. If you plan to culture the cells immediately, swiftly thaw the vial by shaking it in a 37 degrees water bath with clean water and an antimicrobial agent for 40-60 seconds. Remove the vial once a small ice clump persists, ensuring it remains cold. Proceed with all subsequent steps under aseptic conditions. In a sterile flow hood, disinfect the cryovial with 70% ethanol. Then, gently open the vial and transfer the cell suspension into a 15 ml centrifuge tube pre-filled with 8 ml of room temperature culture medium. Gently mix the cells. For cell separation, centrifuge at 300 x g for 3 minutes and dispose of the supernatant. Skipping centrifugation is optional, although any residual freezing medium should be removed after 24 hours. Resuspend the pellet gently in 10 ml of fresh culture medium and divide between two T25 culture flasks. Follow the subculture protocol for subsequent steps.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is rigorously excluded using both PCR-based assays and luminescence-based mycoplasma detection methods. To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

Amelogenin: x,y
CSF1PO: 12
D13S317: 12,14
D16S539: 9,12
D5S818: 11,13
D7S820: 9,10
TH01: 6
TPOX: 8
vWA: 14,19
D3S1358: 16
D21S11: 28,30
D18S51: 12,18
Penta E: 5,15
Penta D: 9,12
D8S1179: 13,14
FGA: 22,25

HT-1080 Cells | 300216

HLA alleles

- A***: 01.01.1900 07:01, 02.01.1900 20:01
- B***: 01.01.1900 03:05
- C***: 02:02:02
- DRB1***: 03:01:01, 04:07:01
- DQA1***: 03:03:01, 05:01:01
- DQB1***: 02:01:01, 03:01:01
- DPB1***: 03:01, 04:01
- E**: 01:01, 01:03