

U-138 MG Cells | 300363

General information

Description	This is one of a number of cell lines derived from malignant gliomas, e.g. U-87-MG, U-118-MG and U-373-MG isolated by J. Ponten and associates from 1966 to 1969. It differs from U-87-MG in morphology and it has a slower proliferation rate. U-138-MG shows strong similarity to U-118-MG, sharing at least six derivative marker chromosomes.
Organism	Human
Tissue	Brain
Disease	Astrocytoma
Synonyms	U-138MG, U-138-MG, U138-MG, U 138 MG, U138MG, U138, 138 MG, 138MG

Characteristics

Age	47 years
Gender	Male
Ethnicity	Caucasian
Morphology	Polygonal
Growth properties	Adherent

Regulatory Data

Citation	U-138 MG (Cytion catalog number 300363)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0020

Biomolecular Data

U-138 MG Cells | 300363

Antigen expression	Blood Type A, Rh+
Isoenzymes	Me-2, 1, PGM1, 1, PGM3, 1, ES-D, 1, AK-1, 1, GLO-1, 1-2, G6PD, B,
Karyotype	Hyperdiploid to pentaploid with several markers, the stemline chromosome number is near triploid with the 2S component occurring at 9.8%. Five markers [t(11,5), t(8q,4), t(19,?18), M1 and M2] were common to most S metaphases. One chromosome 4 could be found in every S metaphase. Chromosome composition was very uniform among cells. Phenotype Frequency Product: 0.0511

Handling

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	1 x 10 ⁴ cells/cm ²
Fluid renewal	2 to 3 times per week
Freeze medium	As a cryopreservation medium, we use 50% basal medium + 40% FBS + 10% DMSO, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

U-138 MG Cells | 300363

**Thawing and
Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

**Incubation
Atmosphere**

37°C, 5% CO₂, humidified atmosphere.

**Shipping
Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

U-138 MG Cells | 300363

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.