Product sheet

CHO-FOLR1 | 305416

[redacted]

Description	[redacted]/[redacted] [redacted] [redacted] [redacted] [redacted] CHO-FOLR1 [redacted] CHO (Chinese Hamster Ovary) [redacted] (FRα) [redacted] FBP [redacted] GPI-anchored [redacted] [redacted] [redacted] FOLR1 [redacted]
Organism	[redacted]
Tissue	[redacted]
Disease	[redacted], [redacted]; [redacted] FOLR1 ([redacted]
Applications	[redacted]; [redacted] FOLR1; [redacted] ADC; [redacted]/[redacted]; [redacted]

[redacted]

Age	[redacted]
Gender	[redacted]
Morphology	[redacted]
Cell type	[redacted]
Growth properties	[redacted]/[redacted]

[redacted]

Citation	CHO-FOLR1 ([redacted] Cytion 305416)
Biosafety level	1
NCBI_TaxID	10029
CellosaurusAccession	CVCL_A8W5

[redacted] CHO-FOLR1 | 305416

GMO Status	GMO-S1: [redacted] CHO [redacted] FOLR1 [redacted]

[redacted]

Receptors expressed	FOLR1 ([redacted] (FRα) [redacted] FBP)

[redacted]

Culture Medium	[redacted]: DMEM:Ham's F12 (1:1), w: 3.1 [redacted]/[redacted] [redacted], w: 2.5 [redacted] L-[redacted], w: 15 [redacted] [redacted]: [redacted] CHO A ([redacted] InSCREENeX; [redacted] InSCREENeX I
Supplements	[redacted]: [redacted] FBS 5% [redacted] Geneticin (G418-Sulfat) [redacted] 0.5 [redacted]/[redacted]
Dissociation Reagent	[redacted]: [redacted]-EDTA
Doubling time	[redacted] 14-16 [redacted]
Subculturing	[redacted]: [redacted] PBS [redacted]
Split ratio	1 [redacted] 5
Seeding density	2 [redacted] 5 x 10^4 [redacted]/[redacted]
Fluid renewal	2 [redacted] 3 [redacted]
Post-Thaw Recovery	[redacted] 1:2 [redacted] 1:3 [redacted] T25 [redacted]
Freeze medium	[redacted] ([redacted] FBS) + 10% DMSO

CHO-FOLR1 | 305416

Thawing and Culturing Cells	1. 2. cryovial -150 3. 37°C 4. 70% 5. 15 6. 300 x g 3 7. 10 8.
Incubation Atmosphere	37°C, 5% $_{CO2}$,
Flask Coating	
Freezing Procedure	
Shipping Conditions	
Storage Conditions	−150 −196 °C

Sterility

PCR