HEK293-CCR8 | 305426

[redacted]

Description	[redacted]/[redacted] [redacted] [redacted] [redacted] [redacted] HEK293-CCR8 [redacted] HEK293 [redacted] [redacted] CCR8 [redacted]
Organism	[redacted]
Tissue	[redacted]
Disease	[redacted]/[redacted]; [redacted] ([redacted] HEK293)
Applications	[redacted]; [redacted] GPCR; [redacted] CCR8; [redacted] CAR-T; [redacted]

[redacted]

Age	[redacted]
Gender	[redacted]
Morphology	[redacted]
Cell type	[redacted]
Growth properties	[redacted], [redacted]

[redacted]

Citation	HEK293-CCR8 ([redacted] Cytion 305426)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_6G18

[redacted] HEK293-CCR8 | 305426

GMO Status	GMO-S1: [redacted] HEK293 [redacted] CCR8 [redacted]

[redacted]

Receptors expressed	CCR8 (CHEMR1 [redacted] CDw198)

[redacted]

Culture Medium	RPMI 1640, w: 2.0 mM [redacted], w: 2.0 [redacted]/[redacted] NaHCO3 ([redacted] Cytion 820700a)
Supplements	[redacted] FBS 10%, [redacted] 1 [redacted], HEPES 10 [redacted], NEAA 1% [redacted]
Dissociation Reagent	[redacted]-[redacted]
Doubling time	[redacted] 24-36 [redacted]
Subculturing	[redacted]
Split ratio	1 [redacted] 5
Seeding density	2 [redacted] 4 x 10^4 [redacted]/[redacted]
Fluid renewal	2 [redacted] 3 [redacted]
Post-Thaw Recovery	[redacted] 1:2 [redacted] 1:3 [redacted] T25 [redacted]
Freeze medium	[redacted] ([redacted] FBS) + 10% DMSO

HEK293-CCR8 | 305426

Thawing and Culturing Cells	1. 2. cryovial -150 3. 37°C 4. 70% 5. 15 6. 300 x g 3 7. 10 8.
Incubation Atmosphere	37°C, 5% $_{CO2}$,
Flask Coating	
Freezing Procedure	
Shipping Conditions	
Storage Conditions	−150 −196 °C

Sterility	PCR