

NCM460 | 305430

Description [REDACTED] NCM460 [REDACTED]
[REDACTED] NCM460 [REDACTED]
NCM460 [REDACTED]

Organism [REDACTED]

Tissue [REDACTED], [REDACTED]

Disease [REDACTED]

Synonyms NCM-460

Age 68 [REDACTED]

Gender [REDACTED]

Ethnicity [REDACTED]

Morphology [REDACTED]

Cell type [REDACTED]

Growth properties [REDACTED]

Citation NCM460 ([REDACTED] Cytion 305430)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_0460

NCM460 | 305430

Tumorigenic

Culture Medium DMEM, w: 4.5 g/l Glucose, w: 4 g/l L-Glutamine, w: 3.7 g/l NaHCO₃, w: 1.0 g/l Penicillin, w: 1.0 g/l Streptomycin

Supplements FBS 10% NEAA 1%

Dissociation Reagent Trypsin

Doubling time 32-38 days

Subculturing PBS, T25

Freeze medium FBS (10%) + 10% DMSO

- Thawing and Culturing Cells**
1. Thaw cryovial in 37°C water bath
 2. Add 10 ml cryovial into 300 x 300 mm cryovial -150 °C
 3. Add 10 ml cryovial into 300 x 300 mm cryovial 37°C
 4. Add 10 ml cryovial into 300 x 300 mm cryovial 70%
 5. Add 10 ml cryovial into 300 x 300 mm cryovial 15
 6. Add 10 ml cryovial into 300 x 300 mm cryovial 3
 7. Add 10 ml cryovial into 300 x 300 mm cryovial 10
 8. Add 10 ml cryovial into 300 x 300 mm cryovial

Incubation Atmosphere 37°C, 5% CO₂

Product sheet

NCM460 | 305430

Flask Coating [REDACTED]

Freezing Procedure [REDACTED]

Shipping Conditions [REDACTED]

Storage Conditions [REDACTED] -150 [REDACTED] -196 °C [REDACTED]

/ / HLA

Sterility [REDACTED] PCR [REDACTED]
[REDACTED] [REDACTED] [REDACTED]