

| 305351

Description	Toledo (DLBCL) (Ig) (transformation)
Organism	
Tissue	
Disease	
Applications	 ,
Synonyms	TOLEDO
Age	
Gender	
Ethnicity	
Morphology	
Growth properties	
Citation	Toledo (Cytion 305351)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_3611

| 305351

[illegible]

| 305351

**Thawing and
Culturing Cells**

1.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
2.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
3.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
4.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
5.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
6.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
7.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.
8.

Remove the cryovial from liquid nitrogen storage and allow it to thaw at room temperature for 1-2 minutes. Do not shake the cryovial. Transfer the cell suspension to a 15 mL conical centrifuge tube. Add 10 mL of pre-warmed cell culture medium. Gently mix the cells by pipetting up and down. Seed the cells into a T75 cell culture flask. Incubate the cells at 37°C, 5% CO₂ in a humidified incubator. After 24 hours, check the cell attachment. If the cells are not attached, repeat the seeding process. Once the cells are attached, replace the medium with fresh medium. The cells should reach confluence within 3-5 days.

**Incubation
Atmosphere**

37°C, 5% CO₂, humidified

**Shipping
Conditions**

Shipped on dry ice, stored at -150 °C to -196 °C, shipped at -150 °C to -196 °C

**Storage
Conditions**

Store at -150 °C to -196 °C, stored at -150 °C to -196 °C

Sterility

PCR negative, endotoxin free, mycoplasma free

Cell suspension is free of mycoplasma, endotoxin, and other contaminants