

# CHO-CXCR4 | 305411MH

## CHO-CXCR4

**Description**

CHO-CXCR4-Medium-high CHO (Chinese Hamster Ovary) CXCR4 CD184

**Organism** CHO

**Tissue** CHO

**Disease** Chinese hamster ovary, non-neoplastic; genetically engineered for CXCR4 surface expression (medium-high expression level)

**Applications** Antibody screening; CXCR4-targeted therapy development; HIV entry research; hematopoietic stem cell biology; flow cytometry

**Synonyms** CHO-CXCR4

## CHO-CXCR4

**Age** CHO

**Gender** CHO

**Morphology** CHO

**Cell type** Epithelial cells

**Growth properties** CHO

## CHO-CXCR4

**Citation** CHO-CXCR4 (Cytion 305411MH)

**Biosafety level** 1

**NCBI\_TaxID** 10029

## CHO-CXCR4 | 305411MH

**CellosaurusAccession** CVCL\_A8W0

**GMO Status** GMO-S1: This CHO derivative contains a construct driving medium-to-high expression of human CXCR4 for GPCR signaling and ligand-binding analyses. This classification applies only within Germany and may differ elsewhere.

**Receptors expressed** CXCR4 (CXCR4)

**Culture Medium** DMEM:Ham's F12 (1:1), w: 3.1 g/l, w: 2.5 g/l L-Glutamine, w: 15 g/l InSCREENeX; InSCREENeX INS-ME-1039

**Supplements** FBS 5% Geneticin (G418-Sulfat) 0.5 mg/ml

**Dissociation Reagent** EDTA

**Doubling time** approx. 14-16 hours

**Subculturing** PBS

**Split ratio** 1 to 5

**Seeding density** 2 to 5 x 10<sup>4</sup> cells/cm<sup>2</sup>

**Fluid renewal** 2-3 times

**Post-Thaw Recovery** 1:2 to 1:3 T25

**Freeze medium** (FBS) + 10% DMSO

**CHO-CXCR4 | 305411MH**

**Thawing and Culturing Cells**

1. Thaw the vial rapidly in a 37°C water bath.
2. Add the contents of the vial to 5 ml of pre-warmed complete medium.
3. Centrifuge at 300 x g for 3 minutes.
4. Resuspend the cell pellet in 10 ml of complete medium.
5. Seed the cells into a T25 flask containing 15 ml of complete medium.
6. Incubate the cells at 37°C, 5% CO<sub>2</sub>, humidified atmosphere.
7. Once cells reach confluence, passage the cells into a T75 flask.
8. Maintain the cells in complete medium.

**Incubation Atmosphere**

37°C, 5% CO<sub>2</sub>, humidified atmosphere.

**Shipping Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

**CHO-CXCR4 / CHO-CXCR4 / HLA**

**Sterility**

CHO-CXCR4 / CHO-CXCR4 / HLA cells are tested for sterility by PCR.