[redacted]4T1-GFP | 305625

[redacted]

Description	4T1-GFP [redacted] 4T1 [redacted] GFP [redacted] 4T1 [redacted] [redacted] 4T1 [redacted] [redacted]
Organism	[redacted]
Tissue	[redacted]
Disease	[redacted]
Metastatic site	[redacted] ([redacted] BALB/c [redacted]); [redacted], [redacted], [redacted]
Applications	[redacted]; [redacted] GFP; [redacted];
Synonyms	4T1-A, 4T1.0, 4T1/WT

[redacted]

Age	[redacted]
Gender	[redacted]
Morphology	[redacted]
Cell type	[redacted]
Growth properties	[redacted]

[redacted]

Citation	4T1-GFP ([redacted] Cytion 305625)
Biosafety level	1

[redacted]4T1-GFP | 305625

NCBI_TaxID	10090
CellosaurusAccession	CVCL_0125
GMO Status	GMO-S1: [redacted] ZsGreen1 [redacted]

[redacted]

Surface antigens	[redacted]
Antigen expression	ZsGreen1 ([redacted])

[redacted]

Culture Medium	RPMI 1640, w: 2.1 mM [redacted], w: 2.0 [redacted]/[redacted] NaHCO3 ([redacted] Cytion 820700a)
Supplements	[redacted] FBS 10%
Dissociation Reagent	[redacted]
Doubling time	12.6 [redacted]
Split ratio	1 [redacted] 3
Seeding density	1 [redacted] 3 x 10^4 [redacted]/[redacted]^2
Fluid renewal	2 [redacted] 3 [redacted]
Freeze medium	[redacted] + 10% DMSO [redacted]

[redacted]4T1-GFP | 305625

Thawing and Culturing Cells	1. [redacted] 2. [redacted] cryovial [redacted] -150 [redacted] 3. [redacted] 37 [redacted] 4. [redacted] 70% 5. [redacted] 15 [redacted] 6. [redacted] 200 x g [redacted] 5 [redacted] 7. [redacted]
Incubation Atmosphere	37°C, 5% $_{CO2}$, [redacted]
Shipping Conditions	[redacted]
Storage Conditions	[redacted] −150 [redacted] −196 °C [redacted]

[redacted]