

XXXXXXXXXXXXXXXXXXXX

XXXXXXXXXXXXXXXXXXXXXXXXXXXX

Product sheet

XXXXXXXXXXHL-1 | 305264

CellosaurusAccession CVCL_0303

XXXXXXXXXXXXXXXXXXXX

Viruses XXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXX40 (SV40)

XXXXXXXXXXXX

Culture Medium XXXXXClaycomb, XXXXXXXX2 mM, FBS 10%, XXXXXXXXXXXXXXX0.1 mM XXXXXXXXXXXX-XXXXXXXXXXXX (XXXXXXXXXXXXXXXXXXXX)
XXXXXXXXXXXXXXXXXXXXXXXXXXXX0.1 mM XXXXXXXXXXXX-XXXXXXXXXXXX
XXXXXXXXXXXXXXXXXXXXXXXXXXXX100 XXXXX= 10 mM XXXXXXXXXXXXXXX
1. XXXXXXXXXXXXXXXL-ascorbic acid 30 mM (50 mL)
2. XXXXXXXXXXXXXXX160 XXX XXXXXXXXXXXXXXXXXXXXXXX(50 XXX) = XXXXXXXXXXXXXXX10 mM
3. XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX0.2 µm XXXXXXXXXXXXXXX
4. XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX100 XXXXXXXXXXXXXXX1 mL XXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXX
5. XXXXXXXXXXXXXXX1 mL XXXXXClaycomb medium 100 mL XXXXXXXXXXXXXXXXXXXXXXX1 XXXXXXXXXXXXXXX

Supplements XXXXXFBS 10%, XXXXXXXX2 mM XXXXXXXXXXXXXXX0.1 mM XXXXXXXXXXXX-XXXXXXXXXXXX

Dissociation Reagent XXXXXXXX

Doubling time XXXXXX24 XXX36 XXXXXXXX

Subculturing XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXPBS XXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXT25 XXX

Split ratio 1 XXX3

Seeding density 2 XXX4 X XXXXXXX/XXX

Fluid renewal XXXXXXX (XXXXXXXXXXClaycomb Medium XXX24 XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX)

Post-Thaw Recovery XXXXXXXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX/XXXXXXXXXX (XXXXXXXXXXXXXXXXXXXX)

DMEM + 10% FBS + 10% DMSC

1.
2. cryovial -150
3. 37°C
4. 70%
5. 15
6. 300 x g 3
7. 10
8.

37°C, 5% CO₂,

25 µg 0.02%

—150 —196 °C

PCR