



Product sheet




SK-LMS-1  | 300125










Antigen expression  Rh+

Isoenzymes Me-2, 2, PGM3, 1-2, PGM1, 1-2, ES-D, 1, AK-1, 1, GLO-1, 1-2, G6PD, B,  0.0027


Tumorigenic  leiomyosarcoma




Karyotype (P12)  3  (+A2, +A3, +C, +D, +E, +F, +G, -A) 






Culture Medium DMEM:Ham's F12 (1:1), w: 3.1  /   w: 2.5  L- w: 15  HEPES, w: 0.5 

Supplements  FBS 10%

Dissociation Reagent 

Subculturing  PBS  T25 
















Split ratio  1:2  1:5


Fluid renewal 2  3 

Freeze medium  ( FBS) + 10% DMSO

SK-LMS-1  | 300125


Thawing and Culturing Cells

1. 
2.  cryovial  -150 
3.  37°C 
4.  70%
5.  15 
6.  300 x g  3 
7.  10 
8. 

Incubation Atmosphere 37°C, 5%_{CO2} 




Flask Coating 

Freezing Procedure 

Shipping Conditions 

Storage Conditions  -150  -196 °C 

 /  / HLA

Sterility  PCR 


SK-LMS-1 **XXXXXXXX** | 300125

XXXXXXXX STR **Amelogenin:** x,y
CSF1PO: 9,1
D13S317: 12
D16S539: 8,11
D5S818: 11,13
D7S820: 8,9
TH01: 6,7
TPOX: 8,9
vWA: 18
D3S1358: 15,16
D21S11: 28,3
D18S51: 14,19
Penta E: 7,13
Penta D: 12,13
D8S1179: 12
FGA: 22,25
PEZ6: ~~XX-XXXXXXXXXX-XXXXXX~~7