Product sheet

HPAF-II | 305088

Description	HPAF-II HPAF-II
Organism	
Tissue	
Disease	
Metastatic site	
Synonyms	HPAF II, HPAFII, HPAF-2, HPAF2, HPAF/CD18, CD18/HPAF, HPAF-II/CD18, CD-18, CD18, CD 18

Age	44
Gender	
Ethnicity	
Morphology	
Growth properties	

Citation	HPAF-II (Cytion 305088)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0313

[redacted] HPAF-II | 305088

[redacted]

[redacted]

Culture Medium	EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO3, w: EBSS ([redacted] Cytion 820100a)
Supplements	[redacted] FBS 10% [redacted] NEAA 1%
Dissociation Reagent	[redacted]
Doubling time	26 [redacted]
Subculturing	[redacted] PBS [redacted] T25 [redacted]
Fluid renewal	2 [redacted] 3 [redacted]
Freeze medium	[redacted] (FBS) + 10% DMSO
Thawing and Culturing Cells	
Incubation Atmosphere	37°C, 5% $_{CO2}$, [redacted]

1. [redacted]
2. [redacted] cryovial [redacted] -150 [redacted]
3. [redacted] 37°C [redacted]
4. [redacted] 70%
5. [redacted] 15 [redacted]
6. [redacted] 300 x g [redacted] 3 [redacted]
7. [redacted] 10 [redacted]
8. [redacted]

HPAF-II | 305088

Flask Coating	
Freezing Procedure	
Shipping Conditions	
Storage Conditions	−150 −196 °C

/ / HLA

Sterility	PCR