Product sheet

[redacted] 3T3-L1 | 400107

[redacted]

Description	[redacted] 3T3-L1 [redacted] 3T3-L1 [redacted] 3T3-L1 [redacted] 3T3-L1 [redacted] 3T3-L1 [redacted] 3T3-L1 [redacted]
Organism	[redacted]
Tissue	[redacted]
Metastatic site	[redacted] ([redacted]; [redacted])
Applications	[redacted] 3T3-L1 [redacted]
Synonyms	3T3 L1, 3T3L1, 3T3-L1 ad, NIH-3T3-L1, NIH3T3-L1

[redacted]

Breed/Subspecies	[redacted]
Age	[redacted]
Gender	[redacted]
Morphology	[redacted]
Cell type	[redacted] / [redacted] ([redacted])
Growth properties	[redacted]

[redacted]

Citation	3T3-L1 ([redacted] Cytion 400107)

3T3-L1 | 400107

Biosafety level	1
NCBI_TaxID	10090
CellosaurusAccession	CVCL_0123
GMO Status	; 3T3-L1 NIH/3T3

Tumorigenic	
Virus susceptibility	, , , , , N-tropic oncornaviruses C
Products	, ,
Ploidy status	
Karyotype	2n 40

Culture Medium	DMEM, w: 4.5 / , w: 4 -, w: 3.7 / NaHCO3, w: 1.0
Supplements	FBS 10%
Dissociation Reagent	
Subculturing	PBS T25
Split ratio	1 5
Seeding density	1 3×10^4 /
Fluid renewal	2 3

3T3-L1 | 400107

Freeze medium	(FBS) + 10% DMSO
Thawing and Culturing Cells	1. 2. cryovial -150 3. 37°C 4. 70% 5. 15 6. 300 x g 3 7. 10 8.
Incubation Atmosphere	37°C, 5% $_{CO2}$,
Flask Coating	
Freezing Procedure	
Shipping Conditions	
Storage Conditions	−150 −196 °C

[redacted]3T3-L1 | 400107

Sterility	[redacted] PCR [redacted] [redacted]