

HEK293-CCR8 Cells | 305426

General information

Description

Disclaimer: The prices displayed for cell lines are exclusively for academic/not-for-profit customers. For commercial entities the price is approximately €6,250.

If you represent a commercial entity or are unsure which category applies, please [contact us](#).

The HEK293-CCR8 cell line is a stable recombinant HEK293 cell line engineered to express the CCR8 receptor at a medium-high level, approximately 22,000 molecules per cell. This cell line was developed using inscreenex's landing pad technology, ensuring precise and reproducible integration of the CCR8 gene at a specific, pre-validated genomic locus. CCR8, also known as CHEMR1 or CDw198, is a G protein-coupled receptor (GPCR) that is predominantly expressed on regulatory T cells (Tregs). It plays a crucial role in the suppression of immune responses within the tumor microenvironment, aiding tumor cells in evading immune detection. Due to its role in maintaining an immunosuppressive environment, CCR8 has become a promising target for novel cancer immunotherapies aimed at enhancing anti-tumor immunity by reducing Treg-mediated suppression.

The expression of CCR8 in this cell line was confirmed using flow cytometry with a target-specific antibody, ensuring consistent and reliable receptor density across the cell population.

Organism

Human

Tissue

Fetal Kidney

Disease

Transformed/immortalized; non-tumorigenic (HEK293 background)

Applications

Immunotherapy research; GPCR pharmacology; CCR8-targeted antibody development; CAR-T cell assays; immuno-oncology screening; Treg biology

Characteristics

Age

Fetus

Gender

Female

Morphology

Epithelial-like

Cell type

Epithelial cells

Growth properties

Monolayer, adherent

Regulatory Data

HEK293-CCR8 Cells | 305426

Citation	HEK293-CCR8 (Cytion catalog number 305426)
----------	--

Biosafety level	1
-----------------	---

NCBI_TaxID	9606
------------	------

CellosaurusAccession	CVCL_6G18
----------------------	-----------

GMO Status	GMO-S1: This HEK293 cell line contains a CCR8 expression construct, enabling chemokine receptor signaling studies. This classification applies only within Germany and may differ elsewhere.
------------	--

Biomolecular Data

Receptors expressed	CCR8 (CHEMR1 or CDw198)
---------------------	-------------------------

Handling

Culture Medium	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
----------------	--

Supplements	Supplement the medium with 10% FBS, 1 mM sodium pyruvate, 10 mM HEPES, 1% NEAA. Add Geneticin (G418-Sulfat) to achieve a final concentration of 1 mg/mL.
-------------	--

Dissociation Reagent	Trypsin-EDTA
----------------------	--------------

Doubling time	approx. 24-36 hours
---------------	---------------------

Subculturing	For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C until the cells detach (5-10 minutes). Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO ₂ , and change the medium every 2-3 days.
--------------	--

Split ratio	1 to 5
-------------	--------

Seeding density	2 to 4 x 10 ⁴ cells/cm ²
-----------------	--

HEK293-CCR8 Cells | 305426

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery

After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere (for adherent cultures) for at least 24 hours.

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

HEK293-CCR8 Cells | 305426

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.