

F81 Cells | 305015

Información general

Description

F81 cells are a feline cell line, specifically derived from feline kidney tissue. These cells are primarily used in research related to the study of feline viruses, including the feline calicivirus (FCV), which is a significant pathogen affecting cats globally. The F81 cell line is noted for its susceptibility to various strains of FCV, making it an indispensable tool in virology research for the development of vaccines and therapeutic strategies aimed at controlling viral infections in felines.

In addition to their use in studying FCV, F81 cells are also utilized in research into other viral pathogens affecting cats, such as feline herpesvirus and feline immunodeficiency virus. Their robustness and ease of culture make F81 cells a valuable model for cellular and molecular biology studies, facilitating advancements in understanding the cellular mechanisms of viral replication and host-pathogen interactions. These cells contribute significantly to translational research, potentially leading to improved diagnostic methods and treatments for feline diseases.

Organism Cat

Tissue Kidney

Synonyms F-81, FK81

Características

Age 3 months

Gender Female

Morphology Endothelial

Growth properties Adherent

Datos normativos

Citation F81 (Cytion catalog number 305015)

Biosafety level 1

NCBI_TaxID 9685

CellosaurusAccession CVCL_9259

F81 Cells | 305015

Datos biomoleculares

Manejo

Culture Medium	EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO ₃ , w: EBSS (Cytion article number 820100a)
Supplements	Supplement the medium with 10% FBS and 1% NEAA
Dissociation Reagent	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Fluid renewal	2 to 3 times per week
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

F81 Cells | 305015

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Control de calidad y análisis molecular

F81 Cells | 305015

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.