

MX-1 Cells | 300296

Información general

Description	The Mx-1 cell line has been established as in vitro culture from the Mx-1 tumor xenograft model of breast carcinoma tissue.
Organism	Human
Tissue	Breast
Disease	Adenocarcinoma, Infiltrating duct carcinoma (IDC)
Synonyms	Mx1, Mxl

Características

Age	29 years
Gender	Female
Ethnicity	Caucasian
Morphology	Epithelial-like
Growth properties	Adherent

Datos normativos

Citation	Mx-1 (Cytion catalog number 300296)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_4774

Datos biomoleculares

Receptors expressed	Estrogen (oestrogen) receptor (-)
----------------------------	-----------------------------------

MX-1 Cells | 300296

Protein expression P53 (-)

Tumorigenic Yes, in nude mice

Manejo

Culture Medium DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Doubling time 30 to 35 hours

Subculturing Remove medium and rinse the adherent cells using PBS without calcium and magnesium (3-5 ml PBS for T25, 5-10ml for T75 cell culture flasks). Add TrypleExpress (1-2ml per T25, 2.5ml per T75 cell culture flask), the cell sheet must be covered completely. Incubate at 37 degree Celsius for 10 minutes. Carefully resuspend the cells, the addition of medium is optional but not necessary, and dispense into new flasks which contain fresh medium. Do not allow the cells to become confluent, subculture once per week. Note: The cells do not form a confluent monolayer. Subculture when a dense layer of cells is observed macroscopically.

Seeding density 2×10^4 cells/cm²

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery Fast

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

MX-1 Cells | 300296

**Thawing and
Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

**Incubation
Atmosphere**

37°C, 5% CO₂, humidified atmosphere.

**Shipping
Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Control de calidad y análisis molecular

MX-1 Cells | 300296

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.