

T2 Cells | 305228

Renseignements généraux

Description

The T2 cell line is a derivative of the T1 human lymphoblastoid cell line and is characterized by its unique properties related to antigen processing and presentation. These cells are deficient in the transporter associated with antigen processing (TAP), resulting in the inability to effectively transport peptides into the endoplasmic reticulum for loading onto major histocompatibility complex (MHC) class I molecules. This deficiency makes T2 cells particularly valuable in immunological research, especially in studies related to the presentation of antigens and the function of MHC class I molecules. By using T2 cells, researchers can better understand the mechanisms of immune recognition and the role of TAP in antigen presentation. T2 cells are also known for their application in cytotoxic T lymphocyte (CTL) assays. Due to their TAP deficiency, these cells express very low levels of surface MHC class I molecules unless exogenous peptides are added. This property allows for the precise study of peptide-MHC interactions and the evaluation of CTL responses to specific antigens. Furthermore, T2 cells are used in vaccine development research, particularly in designing strategies that enhance the presentation of antigens to the immune system. Their unique characteristics make T2 cells a crucial tool in both basic and applied immunology research.

Organism Human

Synonyms T2 (174 x CEM.T2), T2(174 x CEM.T2), 174xCEM.T2, CEMx721.174.T2

Caractéristiques

Morphology Lymphoblast

Growth properties Suspension

Données réglementaires

Citation T2 (Cytion catalog number 305228)

Biosafety level 2

NCBI_TaxID 9606

CellosaurusAccession CVCL_2211

Données biomoléculaires

Manipulation

T2 Cells | 305228

Culture Medium

RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Supplements

Supplement the medium with 10% heat-inactivated FBS

Subculturing

Suspension cells: Remove cells from substrate by pipetting with fresh medium. To obtain single cells, pass the suspension several times through a 22 gauge needle and dispense into new flasks.

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

T2 Cells | 305228

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Contrôle de la qualité et analyse moléculaire

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.