

**A2780-GFP | 305676****Renseignements généraux****Description**

A2780-GFP is a genetically modified derivative of the human ovarian carcinoma cell line A2780, engineered to stably express green fluorescent protein (GFP). The parental A2780 cell line was established from an ovarian tumor in an adult patient and is widely used as a model for epithelial ovarian cancer, particularly in studies of chemotherapy response. It is known for its relative sensitivity to platinum-based agents such as cisplatin, making it a valuable system for investigating mechanisms of drug action and resistance. The GFP-expressing variant preserves these biological characteristics while incorporating a fluorescent reporter for enhanced experimental utility.

The stable expression of GFP enables real-time visualization and quantitative analysis of A2780-GFP cells in a variety of experimental settings. Fluorescence-based detection allows researchers to monitor cell proliferation, morphology, migration, and viability *in vitro*, as well as to track tumor growth and dissemination *in vivo*. This is particularly advantageous in xenograft and metastasis models, where GFP facilitates discrimination of tumor cells from surrounding host tissue. The fluorescent signal is generally stable across passages, although expression levels may vary depending on the transduction method and clonal selection.

A2780-GFP retains the core molecular and phenotypic features of the parental A2780 line, including pathways associated with DNA damage response, apoptosis, and chemotherapeutic sensitivity. As such, it is commonly used in high-content imaging assays, drug screening platforms, and co-culture systems where spatial and temporal resolution of tumor cell behavior is required. The addition of GFP significantly enhances the versatility of this model, supporting applications in ovarian cancer research, therapeutic evaluation, and studies of tumor cell dynamics.

**Organism**

Human

**Tissue**

Ovary

**Disease**

Ovarian endometrioid adenocarcinoma

**Metastatic site**

Primary tumor site (ovary)

**Applications**

Ovarian cancer imaging; GFP-based tumor tracking; *in vivo* fluorescence imaging; combination with parental A2780 for comparative studies; drug efficacy visualization

**Caractéristiques****Age**

Age unspecified

**Gender**

Female

**Ethnicity**

African American

**Morphology**

Epithelial-like

**A2780-GFP | 305676****Cell type** Epithelial cells**Growth properties** Adherent**Données réglementaires****Citation** A2780-GFP (Cytion catalog number 305676)**Biosafety level** 1**NCBI\_TaxID** 9606**CellosaurusAccession** Not assigned (A2780-GFP reporter derivative; parental A2780 CVCL\_1099)**GMO Status** GMO-S1: This human A2780 ovarian cancer line contains a lentiviral GFP expression construct for fluorescence-based imaging. This classification applies only within Germany and may differ elsewhere.**Données biomoléculaires****Antigen expression** GFP**Manipulation****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO<sub>3</sub> (Cytion article number 820700a)**Supplements** Supplement the medium with 10% FBS**Dissociation Reagent** Accutase**Split ratio** 1 to 5**Seeding density** 1 to 3 × 10<sup>4</sup> cells/cm<sup>2</sup>**Fluid renewal** 2 to 3 times per week

**A2780-GFP | 305676**

**Freeze medium**

As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

**Thawing and Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $200 \times g$  for 5 minutes, carefully discard the supernatant containing freezing medium.
7. Follow the procedure described under Post-Thaw Recovery

**Incubation Atmosphere**

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

**Shipping Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

**Storage Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

**Contrôle de la qualité et analyse moléculaire**