

UMR-106 Cells | 305197

Renseignements généraux

Description

UMR-106 is an osteosarcoma cell line derived from a rat model, commonly used in studies investigating bone metabolism, cancer biology, and osteoblast differentiation. These cells are highly responsive to parathyroid hormone (PTH), prostaglandins, and bone-resorbing steroids, making them valuable for research on the regulatory mechanisms of bone cells. The PTH responsiveness of UMR-106 cells is notably greater than that of the related cell line UMR-108, highlighting their unique utility in studies focused on PTH signaling pathways. UMR-106 cells also exhibit the production of alkaline phosphatase, osteocalcin, and other bone-related proteins, which are critical markers in osteoblast research.

In cancer research, UMR-106 cells serve as a model to study the molecular mechanisms underlying osteosarcoma development and progression. They exhibit typical features of cancer cells, such as rapid proliferation and the ability to form tumors *in vivo*, allowing researchers to explore the genetic and epigenetic changes associated with osteosarcoma. These cells are also instrumental in preclinical studies for testing the efficacy and safety of new anti-cancer drugs, providing a reliable system for the preliminary evaluation of therapeutic agents.

Furthermore, UMR-106 cells are utilized to investigate the pathways involved in osteoblast function and differentiation. Researchers have observed that activation of protein kinase C in UMR-106 cells inhibits ATP-induced increases in intracellular calcium levels, providing insights into the complex regulatory networks governing osteoblast activity. The responsiveness of these cells to various stimuli, along with their ability to produce key osteoblastic markers, makes UMR-106 a critical tool in the study of bone biology and the development of strategies to treat bone-related disorders.

Organism	Rat
Tissue	Bone
Disease	Rat osteosarcoma
Synonyms	UMR 106, UMR106

Caractéristiques

Breed/Subspecies	Sprague Dawley
Age	Adult
Morphology	Epithelial
Growth properties	Adherent

Données réglementaires

UMR-106 Cells | 305197

Citation	UMR-106 (Cytion catalog number 305197)
-----------------	--

Biosafety level	1
------------------------	---

NCBI_TaxID	10116
-------------------	-------

CellosaurusAccession	CVCL_3617
-----------------------------	-----------

Données biomoléculaires

Receptors expressed	Parathyroid hormone(PTH), 1-25(OH)2D3(bone resorbing steroid hormone)
----------------------------	---

Manipulation

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

Supplements	Supplement the medium with 10% FBS
--------------------	------------------------------------

Dissociation Reagent	Accutase
-----------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	---

UMR-106 Cells | 305197

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Contrôle de la qualité et analyse moléculaire

UMR-106 Cells | 305197

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.