

## VERO Cells | 605372

## Renseignements généraux

**Description**

VERO cells are widely used in developing vaccines, in the study of viral infections or malaria, and in tumor immunology and immunotherapy studies. VERO cells were derived from the kidney of an African green monkey in the 1960s by a group of Japanese scientists at Chiba University in Japan.

One of the critical characteristics of VERO cells is their rapid growth rate, with a population doubling time of approximately 24 hours. This, combined with their stability and high viral titers, makes them an ideal choice for vaccine production. As a prominent example, a Vero cell-derived vaccine for Japanese encephalitis is widely used and licensed in many countries worldwide.

Vero cells were pivotal in the development of vaccines for a plethora of infectious diseases, including the rubella virus, Ross River virus, herpes simplex virus, measles virus, and poliovirus. Vero cells are renowned for their capacity for virus production, growth, and maintenance under optimized culture conditions, making them an invaluable resource in viral vaccine production. The role of Vero cells extends to the generation of viral vectors, crucial for both vaccine development and tissue engineering applications, and virus isolation.

Different VERO cell lines, such as Vero 76 and the subclone Vero E6, offer unique characteristics suited to various research and production needs. Vero 76 cells are known for their robust growth and are widely used in vaccine production due to their high virus yield capabilities. Vero E6, on the other hand, exhibits specific properties that make it particularly useful for studying certain viruses, including enhanced sensitivity to the Ebola virus and SARS-CoV-2. This subclone's unique interaction with viruses makes it valuable for viral pathogenesis studies and antiviral drug screening.

**Organism** Chlorocebus sabaeus (Green monkey)

**Tissue** Kidney

**Applications** Transfection host

**Synonyms** Vero, VeroCCL81, Vero 81, Verda reno

## Caractéristiques

**Age** Adult

**Gender** Female

**Morphology** Epithelial-like

**Growth properties** Monolayer, adherent

## Données réglementaires

## VERO Cells | 605372

**Citation** VERO (Cytion catalog number 605372)

**Biosafety level** 1

**NCBI\_TaxID** 60711

**CellosaurusAccession** CVCL\_0059

## Données biomoléculaires

**Receptors expressed** Despite not being interferon deficient, VERO cell line possesses the interferon-alpha/beta receptor, allowing them to respond normally when recombinant interferon is added to their culture medium.

**Viruses** Verotoxin detection of virus in ground beef

**Virus susceptibility** Poliovirus 1, 2, 3, Getah, Ndumu, Pixuna, Ross River, Semliki Forest, Paramaribo, Kokobera, Modoc, Murutucu, Germiston, Guaroa, Pongola, Tacaribe, SV-5, SV40, rubeola, rubellavirus, reovirus 1, 2, 3, simian adenoviruses

**Reverse transcriptase** Negative

**Mutational profile** Vero cells have a homozygous 9-Mb deletion on chromosome 12 that results in loss of the type I interferon gene cluster and the cyclin-dependent kinase inhibitors CDKN2A and CDKN2B.

## Manipulation

**Culture Medium** DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO<sub>3</sub> (Cytion article number 820400a)

**Supplements** Supplement the medium with 10% FBS

**Dissociation Reagent** Accutase

**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

**Seeding density**  $1 \times 10^4$  cells/cm<sup>2</sup>

## VERO Cells | 605372

**Fluid renewal** 2 to 3 times per week

**Freeze medium**

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

**Thawing and Culturing Cells**

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

**Incubation Atmosphere**

37°C, 5% CO<sub>2</sub>, humidified atmosphere.

**Shipping Conditions**

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

## VERO Cells | 605372

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196$  °C. Storage at  $-80$  °C is acceptable only as a short interim step before transfer to liquid nitrogen.

## Contrôle de la qualité et analyse moléculaire

### Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.